arXiv:1301.3602v1 [math.ST] 16 Jan 2013

FOURIER TRANSFORM METHODS FOR PATHWISE COVARIANCE ESTIMATION IN THE PRESENCE OF JUMPS

CHRISTA CUCHIERO AND JOSEF TEICHMANN

Abstract. We modify (classical) jump-robust estimators of integrated realized covariance to estimate the Fourier coefficients of the instantaneous stochastic covariance. By means of a central limit theorem for those Fourier coefficient estimators we are then able to prove consistency and a (pointwise) central limit theorem for the non-parametrically reconstructed instantaneous covariance process itself. The procedure is – by methods of Fourier analysis – robust enough to allow for an iteration and we can therefore show theoretically and empirically how to estimate the integrated realized covariance of the instantaneous stochastic covariance process. We also explain a surprising shrinkage phenomenon for the constructed Fourier estimators, i.e., in comparison to classical (local) estimators of instantaneous variance the asymptotic estimator variance of the Fourier estimator is smaller by a factor 2/3, but an additional (pointwise small) bias appears. We apply these techniques to robust calibration problems for multivariate modeling in finance, i.e. the selection of a pricing measure by using time series and derivatives' price information simultaneously. "Robust" here means that re-calibration is more stable over time, that the estimation procedures of, e.g., instantaneous covariance also work in the presence of jumps, and that the procedures are as robust as possible with respect to input deficiencies.

1. Robust Calibration

The recent difficulties in the banking and insurance industry are to some extent due to insufficient modeling of multivariate stochastic phenomena which appear in financial markets. There are several reasons why modeling is insufficient, but the two most important ones are the following: first, realistic multivariate models are difficult to calibrate to market information due to a lack of analytic tractability, hence oversimplified models are in use in delicate multivariate situations, and, second, usually either time series data or derivatives' prices are used to select a model from a given model class but *not both* sorts of available information simultaneously. We do not speak here about phenomena which should not be considered stochastic and where stochastic modeling is inappropriate at all, but we concentrate on multivariate time series of liquid instruments, where appropriate and easy to implement stochastic modeling is needed. Applications are risk management, pricing and hedging of instruments, and portfolio management.

2010 *Mathematics Subject Classification.* 60F05, 60G48.

Key words and phrases. non-parametric variance estimation, Fourier analysis, jump-diffusion, jump-robust estimation techniques.

The first author gratefully acknowledges the partial financial support from the European Research Council (ERC, grant FA506041). The first and second author gratefully acknowledge the support from the ETH-foundation.

We aim to develop methods which allow for *robust calibration*, i.e., estimation and calibration of a model in a well specified sense simultaneously from time series and derivatives' prices data in order to select *a pricing measure*. It is useful to demonstrate what we actually mean with *robust calibration* by means of an example: take a Heston model with possibly state-dependent jumps in the log-price process and variance

$$dX_t = \kappa(\theta - X_t)dt + \sigma\sqrt{X_t}dW_t + dL^1_t,$$
$$dY_t = \left(\mu - \frac{X_t}{2}\right)dt + \sqrt{X_t}dB_t + dL^2_t,$$

where X is the stochastic variance process and Y the log-price of a stock. L^1 and L^2 denote the possibly state-dependent jump parts. Through robust calibration we have to identify the initial value X_0, Y_0, the five parameters $\mu, \kappa, \theta, \sigma$ and the correlation between the Brownian motions ρ, and the compensators of the jump parts in order to specify the model for purposes of pricing, hedging or risk management.

Apparently at least the initial values X_0, Y_0, and the parameters σ and ρ do not change under equivalent measure changes, so in principal the parameters X_0, Y_0, σ, ρ can be identified from the observation of a single trajectory. On the other hand market implied values for those parameters should coincide with values estimated from the time series if the model is close to correct. Here "market implied values" means to choose model parameter values such that the model's derivatives' prices and the market prices coincide as well as possible. From basic theory of mathematical finance we know that the calculation of derivatives' prices has to be done under an equivalent martingale measure for the process $\exp(Y)$, but X_0, Y_0, σ, ρ do not change under equivalent measure changes.

Formally speaking we have defined the above model on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and we consider equivalent measures $\mathbb{Q} \sim \mathbb{P}$. Having specified a set of parameters Θ and a semimartingale S^θ depending on parameters $\theta \in \Theta$, we can then define an equivalence relationship, namely $\theta_1 \sim \theta_2$ if $S^{\theta_1}_* \mathbb{P} = S^{\theta_2}_* \mathbb{Q}$, i.e., equality for the respective measures on the canonical probability space of càdlàg paths, for some $\mathbb{Q} \sim \mathbb{P}$. This equivalence relation defines orbits on Θ and the space of orbits $\Theta/\sim$, where the latter set is the set of invariant parameters, i.e., those parameters of S^θ, which are not touched by equivalent measure changes. For sufficiently rich parametric models this set is usually not trivial.

Having this basic stochastic fact in mind, it should be clear that non-parametric estimation of instantaneous covariance processes is *the* important task to be performed, since it yields – in the previous concrete case – information on σ, ρ and the trajectory of instantaneous stochastic variance $t \to X_t$ along the observation interval. The parameters κ and θ, the characteristics of the jumps will rather be calibrated from derivatives' prices (except the Blumenthal-Getoor index), since they cannot be identified from time series information. The difference between the martingale measure identified through calibration and the real world measure is most delicate to estimate from the time series information: we do not comment on this issue here.

To be precise on time series data: we think of intraday price data for liquid instruments along periods of months up to years, such that we have about 10^5 data points available without loosing assumptions on time-homogeneity.

We call a calibration procedure *robust*, if it follows the subsequent steps:

(i) estimate parameters which are invariant under equivalent measure changes from time series by econometric means together with respective confidence sets.
(ii) calibrate the partially specified model via a non-linear (robust) pricing operator to market prices and determine the rest of the parameters. Here we do neither assume *one* model anymore, nor do we expect the pricing operator to provide *one price*, compare to the recent work [3].
(iii) reject the model if the second step calibration is producing time-inconsistent or time-dependent results when monitored along a time series of derivatives' prices.

There is also another reason why time series methods should enter the field of multivariate model selection in mathematical finance. In fact, due to the high dimensional parameter space and the lack of liquidly traded options whose payoffs depend on several underlyings, calibration is particularly challenging. This difficulty can be tackled to a certain extent by additionally using time series of asset prices, from which – according to the above calibration principle – those parameters which remain invariant under equivalent measure changes can be determined.

Based on this concept, the goal of the present article is to find methods which allow to estimate *non-parametrically* the instantaneous stochastic variance of the instantaneous stochastic variance process X, and, to estimate instantaneous stochastic correlation between the log-price process Y and the stochastic variance process X. This involves a two step procedure where we first need to recover the realized path of the instantaneous variance, from which we can then – by iteration of the estimator – estimate the second order quantities.

In the literature the problem of identification of a pricing measure under time series information is usually solved by setting up high-dimensional filtering problems. From the filtering point of view our method can be understood as choosing – by direct non-parametric estimation techniques – excellent priors concentrated around invariant parameters in order to initialize the filtering procedure.

2. A REMARK ON PATHWISE ESTIMATION PROCEDURES

Robust calibration refers to splitting the task of model identification into an estimation and a calibration subtask. This leads to the necessity of finding pathwise estimation procedures for, e.g., instantaneous covariance processes, where we look for robustness rather in the statistical sense. More precisely the presence of non-equidistant and non-equal grids for different log-price processes should not influence the procedures, i.e., we have to encode market information into quantities which depend in *continuous ways* on the observations. This is the topic of this second introductory section, where we argue why to apply Fourier estimation techniques as outlined, e.g., in [9]. We are aware that more classical estimation methods for instantaneous covariance could also be applied for our purposes, but we try to argue that Fourier techniques might appear more robust in the light of the following arguments.

We consider multivariate stochastic models to describe the joint stochastic evolution of several price processes for purposes of pricing, hedging and forecasting. Such models typically incorporate

- a stochastic covariance process, yielding stochastic volatility of each single asset and stochastic correlation between the different assets, and

- jumps, whose jump measures can exhibit a rich dependence structure.

We assume that we discretely observe a path of the log-price process with moderately high frequency (up to minutes) and we would like to estimate from this path stochastic covariance as well as possible. Since the problem is multivariate we should also care about the following two problems:

- the observation grid might be not equidistant.
- observation grids of different prices might differ.

This amounts to develop a theory which has an additional robustness, or in other words, it amounts to use estimators where the gridding plays a minor role. Both aspects are discussed in this introductory section from an basis expansion point of view.

The estimation of *invariant parameters* from discrete observations of a path of a semimartingale Y on a finite interval $[0,T]$ is related to the question how to represent this information in a *robust way*, which in this context simply means *as continuous as possible*. We would like to take a particular point of view on this question, which allows to unify several approaches present in the literature. Consider $L^2([0,T])$ the Hilbert space of complex valued, square-integrable functions on the interval $[0,T]$ and a finite orthonormal system $(e_1,\ldots,e_n)$ in it. The orthogonal projection of an element $f\in L^2([0,T])$ onto the span of $(e_1,\ldots,e_n)$ can then be represented as $\pi_n(f)=\sum_{i=1}^n e_i\,\langle f,e_i\rangle$, hence the calculation of the coordinates $\langle f,e_i\rangle$, for $i=1,\ldots,n$ is key if we want to calculate the projection $\pi_n(f)$. The coordinates are continuous functions of the full paths with respect to the L^2- and even to the L^∞-topology, but continuity is more delicate an issue when it comes to coefficients of the "increment" process, i.e., when we aim to represent the derivative of f

$$\pi_n(f')=\sum_{i=1}^n e_i \int_0^T e_i(s)df(s)\,.$$

Here apparently continuity of the coordinates is not generically true (since, e.g., not every element $f\in L^2([0,T])$ will be differentiable), but remains true if integration by parts works, for instance if $e_i\in C^1$, for $i=1,\ldots,n$. The representation of the derivative f' is important in our setting since this corresponds to the representation of the increments of a stochastic process Y.

Let us assume now that the path of the semimartingale Y is observed on time points $0=t_0^n<t_1^n<\ldots<t_n^n=T$: this set of time points can be encoded in the ("discrete observation") orthonormal system $\varepsilon_i^n=\frac{1}{\sqrt{t_i^n-t_{i-1}^n}}1_{[t_{i-1}^n,t_i^n]}$ for $i=1,\ldots,n$. Then the discrete observation (i.e., observation together with their time points of observation) is bijectively encoded through the following sum of stochastic integrals

$$\sum_{i=1}^n (Y_{t_i^n}-Y_{t_{i-1}^n})\frac{1}{t_i^n-t_{i-1}^n}1_{[t_{i-1}^n,t_i^n]}=\sum_{i=1}^n \varepsilon_i^n\int_0^T \varepsilon_i^n(s)dY_s$$

if tacitly $Y_0:=0$ is assumed. Since many estimators will be formulated as functions of increments, we have to understand the previous expression's continuity properties. In particular this means that proper encoding in the case of discrete observations should rather be formulated in terms of stochastic integrals with respect to the path than in terms of integrals along the path. However, stochastic integrals are known to be rather measurable than continuous for generic integrands on path spaces. Only when the basis functions satisfy some differentiability properties allowing for integration by parts, i.e., for $e \in C^1([0,T])$

$$\int_0^T e(s) dY_s = Y_T e(T) - \int_0^T Y_s e'(s) ds \,,$$

the coordinates $\int_0^T e_i(s) dY_s$ are *continuous* with respect to the path on path space.

It might therefore be wise

- to formulate the theory in a coordinate-free way, i.e., estimators can have inputs coming from different orthonormal systems.
- to allow a change of coordinates for the estimated quantities, i.e., when the input is given on a discrete grid (which corresponds to a basis decomposition with respect to $(\varepsilon_i^n)_{1 \leq i \leq n}$), then output could be produced with respect to any other reasonable orthonormal system.

We shall not fully follow this program, since we do start with discrete observations, i.e., coordinates with respect to $(\varepsilon_i^n)_{1 \leq i \leq n}$, but we shall formulate the theory such that we can estimate consistently, together with a central limit theorem, coordinates $\langle e_i, X \rangle$ for $i = 1, \dots, n$ with respect to any chosen orthonormal system in $L^2([0,T])$. In [8] additionally all estimators are written in terms of Fourier-coordinates $\langle e_i, X \rangle$ for $i = 1, \dots, n$, which has several procedural advantages with respect to non-equidistant or non-equal gridding. We insist that we could formulate our theoretical findings also in this spirit, but in this article we start by writing estimators functional on increments. The main idea now is the following: classical estimators of the form

$$V^n(X) = \sum_{i=1}^n (t_i^n - t_{i-1}^n) g\left(\frac{Y_{t_i^n} - Y_{t_{i-1}^n}}{\sqrt{t_i^n - t_{i-1}^n}} \right)$$

will play a major role, where – under continuity assumptions on g – continuous dependence on discrete observations is clear. On refining grids we shall often obtain consistency, i.e.,

$$V^n(X) \to \int_0^T \rho_g(X_s) ds$$

for some function ρ_g in a consistent way, possibly together with a central limit theorem. The announced "change of coordinates" is then performed through the following sequence of estimators

$$V_j^n(X) = \sum_{i=1}^n (t_i^n - t_{i-1}^n) e_j(t_i^n) g\left(\frac{Y_{t_i^n} - Y_{t_{i-1}^n}}{\sqrt{t_i^n - t_{i-1}^n}} \right), \tag{2.1}$$

for $j = 1, \dots, N$, which converge to

$$\int_0^T e_j(s) \rho_g(X_s) ds \,,$$

for $j = 1, \dots, N$ in a consistent way together with a central limit theorem. This new interpretation of estimators for integrated quantities allows to obtain estimators for coordinates of the spot quantity with respect to any other orthogonal system $(e_i)_{i=1,\dots,N}$.

In view of robustness, which means continuity here, we choose a coordinate system whose coordinates depend as continuous as possible on the observation path $t \mapsto \rho_g(X_t)$ in order to smoothen input deficiencies. This advantage will be important for the estimation of quantities related to X as we shall do it at the end of this work. In particular we believe that the change of coordinates from the orthogonal system $(\varepsilon_i^N)_{1\leq i\leq N}$ to another, possibly more smoothing system like the Fourier basis, $(e_i)_{i=1,\ldots,N}$, can be of advantage when it comes to quality of approximation. Notice also that V_j^n with respect to the basis $(\varepsilon_i^N)_{1\leq i\leq N}$ (where the basis elements are given on a coarser grid corresponding to N) would correspond just to the well-known local realized variance estimators on sliding, non-overlapping windows.

Having constructed estimators for coordinates with respect to a basis $(e_i)_{i=1,\ldots,N}$, we know the projection of the path on the space spanned by $(e_i)_{i=1,\ldots,N}$, but since every coordinate comes with an estimation error, we have to understand how those errors influence the reconstruction. This is the very reason why we introduced another cardinality $N \leq n$ of the (reconstruction) orthonormal system. The choice of N amounts to understanding the size of sums with random error coefficients $\mathcal{E}_i$

$$\sum_{i=1}^{N} e_i \, \mathcal{E}_i \, ,$$

which is small in the L^2-sense on $[0,T]$ and on the observation probability space if $\sum_{i=1}^N \mathbb{E}[\mathcal{E}_i^2]$ is small. The size of this sum determining the L^2-error will depend on the variance of $\mathcal{E}_i$ and on the number N. We see here that – even though we would like to take $N \leq n$ as large as possible – the accuracy of reconstruction is bounded by the size of the previously described error, which grows with N.

If we ask for the L^∞-error in reconstructing the path we should rather consider $t \mapsto \sum_{i=1}^N e_i(t) a_i \, \mathcal{E}_i$, where a_i are an additional weighting sequence optimizing the L^∞-approximation (e.g., Féjer weights in case of the Fourier basis). This leads to an error random variable with expectation

$$\sum_{i=1}^{N} e_i(t) a_i \, \mathbb{E}\left[\mathcal{E}_i\right]$$

and second moment

$$\sum_{i,j=1}^{N} e_i(t) e_j(t) a_i a_j \, \mathbb{E}\left[\mathcal{E}_i \mathcal{E}_j\right] \, ,$$

for $t \in [0,T]$. Assuming for a moment that the errors for the estimation of different coordinates are i.i.d. with vanishing expectation, then the estimator variance equals

$$\mathrm{Var}\left[\mathcal{E}_1\right] \sum_{i=1}^{N} \left(e_i(t)\right)^2 a_i^2 \, ,$$

for $t \in [0,T]$. This error quantity depends on the particular choice of the orthonormal system and the structure of the errors; its size might differ considerably. In particular very localized orthonormal systems rather lead to truncated sums of errors, such that "diversification" effects do not take place. We therefore believe that orthonormal systems like the Fourier basis might bring better results in some cases.

Another aspect for the choice of orthonormal systems besides *continuity of representation of path information* and *control of L^∞-errors* is related to *locality of the estimator*: when we ask for an estimated value of $\rho_g(X_t)$ at one point in time t, we can use in principle all the information along the observed path to infer that value. How information is weighted as a function of time distance to t is strongly influenced by the choice of the basis. Local estimators, like most of the classical estimators, use a comparably small sliding window to estimate the instantaneous quantity $\rho_g(X_t)$, where other, non-local estimators use exponential weights, or algebraically decreasing weights of information along the path.

The Fourier-Féjer estimator, which we introduce in the sequel, satisfies all outlined requirements: we have continuity on path space of functionals $Y \mapsto \int e_i(s)\, dY_s$. Error control might lead to "diversification effects", since sums are never truncated, and the estimators are non-local, i.e., the information of the whole time series enters into estimators of instantaneous covariance. We therefore expect that the estimator variance of Fourier-Féjer estimators might be smaller than for estimators written with respect to the basis $(\varepsilon_i^N)_{1 \leq i \leq N}$, which is indeed the case as proved in Theorem 6.6, see also Remark 6.7. On the other hand an additional bias will (has to) appear, which, however, is pointwise quite small.

This shrinkage phenomenon is confirmed by Figure 1 below, which shows a comparison between the classical local realized variance estimator and the Fourier-Féjer estimator. In particular, the variance of the Fourier-Féjer estimator is comparable with the one of a James-Stein shrinkage variant of the classical estimator.

In our illustration example the underlying semimartingale Y is a drifted Brownian motion with constant variance, that is,

$$dY_t = b_t^Y\, dt + \sqrt{X} dZ_t,$$

where b^Y denotes the drift, Z a standard Brownian motion and X the deterministic constant variance, which we aim to measure on a coarser grid given discrete observations of Y. The Fourier-Féjer estimator $\widehat{X}_t^{n,N}$ is described in detail in the following sections and given by (6.2) (here applied with $g(y) = y^2$ and $T = 1$). As we shall prove in the sequel it is asymptotically normal $\mathcal{N}(X, \frac{4X^2}{3}\frac{N}{n})$ as $n, N \to \infty$ and $\lim \frac{n}{N^\gamma} = K$ for some constant K and $\gamma > 1$. In comparison, the classical estimator is given by

$$\widehat{\widehat{X}}_t^{n,N} = \sum_{j=1}^{N} \varepsilon_j^N(t) \sum_{i=1}^{n} (Y_{t_i^n} - Y_{t_{i-1}^n})^2 \varepsilon_j^N(t_{i-1}^n)$$

with $\varepsilon_j^N(t) = 1_{[t_{j-1}^N, t_j^N]}(t) \frac{1}{\sqrt{t_j^N - t_{j-1}^N}}$, $t_j^N = \frac{j}{N}$ for $j = 1, \ldots, N$ and $t_i^n = \frac{i}{n}$ for $i = 1, \ldots, n$, which corresponds to

$$\sum_{j=1}^{N} \varepsilon_j^N(t) V_j^n(X),$$

where $V_j^n(X)$ is given in (2.1) with $e_j = \varepsilon_j^N$ and $g(y) = y^2$. According to [5, Theorem 13.3.3], $\widehat{\widehat{X}}_t^{n,N}$ is asymptotically normal $\mathcal{N}(X, 2X^2\frac{N}{n})$ as $n, N \to \infty$ and $\lim \frac{n}{N^\gamma} = K$ for some constant K and $\gamma > 1$. In particular, the variance of the Fourier-Féjer estimator is $\frac{2}{3}$ of the one of the classical estimator. A similar variance reduction phenomenon can be achieved by applying the James-Stein estimator to

$\widehat{\widehat{X}}_t^{n,N}$ and considering the following shrinkage estimator:

$$\widehat{\widehat{\widehat{X}}}_t^{n,N} = \left(1 - \frac{(M-2)2X^2N}{n\left\|\widehat{\widehat{X}}^{n,N}\right\|^2}\right)\left(\widehat{\widehat{X}}_t^{n,N} - \frac{1}{M}\sum_k \widehat{\widehat{X}}_{t_k}^{n,N}\right) + \frac{1}{M}\sum_k \widehat{\widehat{X}}_{t_k}^{n,N},$$

where M denotes the number of evaluation points of $\widehat{\widehat{X}}^{n,N}$.



Figure 1. Comparison of estimators

The remainder of the article is organized as follows. In Section 3 we introduce the assumptions on the log-price and the instantaneous covariance process and Section 4 gives an overview of the different steps in our estimation procedure. Section 5 is dedicated to jump robust estimators for the Fourier coefficients, while in Section 6 consistency and a central limit theorem are proved for the Fourier-Féjer instantanous covariance estimator. In Section 7 we consider estimators for the integrated covariance of the instantaneous covariance process and Section 8 concludes with simulation results.

3. SETTING AND ASSUMPTIONS

Throughout we work on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0\leq t\leq T}, \mathbb{P})$, where we consider a d-dimensional (discounted) asset price process $(S_t)_{0\leq t\leq T}$, which is supposed to be nonnegative, and adapted to the filtration $(\mathcal{F}_t)_{0\leq t\leq T}$. Due to the non-negativity of S we further assume

$$S_t = (\exp(Y_{t,1}), \ldots, \exp(Y_{t,d}))^\top, \quad 0 \leq t \leq T,$$

where $(Y_t)_{0\leq t\leq T}$ denotes the d-dimensional (discounted) logarithmic price process starting at $Y_0 = y \in \mathbb{R}^d$ a.s. Due to no-arbitrage consideration S and thus also Y are supposed to be semimartingales with a rich structure of jumps.

Furthermore, let us introduce some mild structural assumptions on the log-price process Y, namely that it is an Itô-semimartingale of the following form:

Assumption (H)**.** *The logarithmic price process Y satisfies*

$$\begin{aligned} Y_t = y &+ \int_0^t b_s^Y ds + \int_0^t \sqrt{X_{s-}} dZ_s + \int_0^t \int_{\mathbb{R}^d} \chi(\xi)(\mu^Y(d\xi, ds) - K_s(d\xi)ds) \\ &+ \int_0^t \int_{\mathbb{R}^d} (\xi - \chi(\xi))\mu^Y(d\xi, ds), \end{aligned} \tag{3.1}$$

where $\sqrt{X}$ denotes the unique matrix square root on S_d^+, the space of positive semi-definite matrices, and

- *Z is a d-dimensional Brownian motion,*
- *b^Y an $\mathbb{R}^d$-valued locally bounded process,*
- *X a càdlàg process taking values in S_d^+ and*
- *$\mu^Y(d\xi, dt)$ the random measure associated with the jumps of Y, whose compensator is given by $K_t(d\xi)dt$, where $K_t(d\xi) = K_t(\omega, d\xi)$ is for each (ω, t) a measure on $\mathbb{R}^d$.*

Remark 3.1. (i) Usually, e.g., in [5], the assumption of an Itô-semimartingale is formulated in terms of the *Gringelions representation*, which means that there exists an extension of the probability space, on which are defined a d'-dimensional Brownian motion W and a Poisson random measure $\mathfrak{p}$ with Lévy measure λ such that

$$\begin{aligned} Y_t = y &+ \int_0^t b_s^Y ds + \int_0^t \sigma_s dW_s + \int_0^t \int_{\mathbb{R}^d} \delta(\xi, s-) 1_{\{\|\delta\| \leq 1\}} (\mathfrak{p}(d\xi, ds) - \lambda(d\xi)ds) \\ &+ \int_0^t \int_{\mathbb{R}^d} \delta(\xi, s-) 1_{\{\|\delta\| > 1\}} \mathfrak{p}(d\xi, ds), \end{aligned}$$

where σ_t is an $\mathbb{R}^{d \times d'}$-valued predictable process such that $\sigma^\top \sigma = X$ and δ is a predictable $\mathbb{R}^d$-valued function on $\Omega \times \mathbb{R}^d \times \mathbb{R}_+$. In view of applications to (affine) processes, whose characteristics are given in terms of representation (3.1), we prefer to use the formulation of Assumption (H) and do not use the Gringelions representation, since it involves an extension of the probability space and the specific form of δ and $\mathfrak{p}$ is not evident.

(ii) The conditions on the characteristics in Assumption (H) correspond to those of [5, Assumption 4.4.2 (or (H))] with the only difference that for the moment we do not assume some kind of local integrability on the jump measure, as in (iii) of [5, Assumption 4.4.2].

Note that the assumption of an Itô-semimartingale is satisfied by all continuous-time models used in mathematical finance. Indeed, beside the assumption of *absolutely continuous* characteristics, this is the most general model-free setting which is in accordance with the no-arbitrage paradigm. This assumption is actually also the only one needed to prove consistency of the Fourier-Féjer instantaneous covariance estimator, denoted by $\widehat{X}_t$. However for establishing a central limit theorem we also need some structural assumptions on the instantaneous covariance process X:

Assumption (H1)**.** *Assumption* (H) *holds and the instantaneous covariance process* X *is an Itô-semimartingale of the form*

$$X_t = x + \int_0^t b_s^X ds + \sum_{j=1}^{p} \int_0^t Q_s^j dB_{s,j} + \int_0^t \int_{S_d} \xi \mu^X(d\xi, ds)$$

where

- B *is* p*-dimensional Brownian motion, which can be correlated with* Z*, the Brownian motion driving the log-price process, such that* $d\langle Z_i, B_j\rangle_t = \rho_{t,ij}dt$*, where* ρ_{ij} *is adapted càglàd for all* $i \in \{1, \ldots, d\}$ *and* $j \in \{1, \ldots, p\}$*,*
- b^X *is an* $\mathbb{R}^{d\times d}$*-valued locally bounded predictable process,*
- $(Q^j)_{j\in\{1,\ldots,p\}}$ *is an adapted càglàd process taking values in* S_d*,*
- μ^X *is the random measure associated with the jumps of* X*, whose compensator is given by* $F_t(d\xi)dt$*, where* $F_t(d\xi) = F_t(\omega, d\xi)$ *is for each* (ω, t) *a measure on* S_d *such that the process*

$$\left(\int_{S_d} (\sqrt{X_{t-} + \xi} - \sqrt{X_{t-}})F_t(d\xi)\right)_{t\geq 0} \tag{3.2}$$

is locally bounded.

Moreover, both processes X_t *and* X_{t-} *take their values in* S_d^{++}*, the set of all (strictly) positive semidefinite* $d \times d$ *matrices. Furthermore, the drift process* b^Y *of* Y *is additionally assumed to be adapted and càglàd.*

Remark 3.2. (i) Assumption ($H1$) corresponds essentially to [5, Assumption 4.4.3 (or (K))] or [1, Assumption ($H1$)], respectively. The main difference is that we require X to be an Itô-semimartingale, whereas in the above references this condition is explicitly stated for $\sqrt{X}$. Since we assume additionally that X_t and X_{t-} take values in S_d^{++}, $\sqrt{X}$ is again an Itô-semimartingale. Local boundedness then also holds for the drift and for the compensator of the jumps of $\sqrt{X}$, where the latter is a consequence of condition (3.2). The motivation to state Assumption ($H1$) in terms of X stems again from applications to S_d^+-valued affine processes, where the characteristics of $\sqrt{X}$ would have a much more complicated form than the simple affine dependence on X.

(ii) Also in view of affine processes we prefer the formulation in terms of a Brownian motion B, correlated with Z instead of decomposing B into Z and another independent Brownian motion.

(iii) Concerning the jump part we implicitly assume that the jumps of X are of finite variation, whence we can avoid the introduction of a truncation function. This is not restrictive in our case, since in the central limit theorem below continuity of $t \mapsto X_t$ is needed.

4. The role of pathwise covariance estimation in robust calibration

In this article we combine jump robust estimators (already considered for integrated covariance estimation) with instantaneous covariance estimation based on Fourier methods, as introduced by Malliavin and Mancino [8, 9]. More precisely, our method consists in conducting the following three steps:

(1) the first step is to reconstruct *non-parametrically* from *discrete observations* along an equidistant time grid $\Lambda_1 = \{t_0^n < \ldots < t_{\lfloor nT\rfloor}^n = T\}$ with step

width $\frac{1}{n}$ of the log-price process Y a discrete set of points on the trajectory of the instantaneous covariance process X along a coarser, equidistant time grid $\Lambda_2 \subset \Lambda_1$:

$$(Y_t)_{t\in\Lambda_1} \longrightarrow (\widehat{X}_t)_{t\in\Lambda_2}\,.$$

(2) using then the reconstructed path $\widehat{X}$ along the coarser grid Λ_2 allows in principle to repeat the first step, or at least to estimate integrated quantities of that discretely given trajectory, for instance to compute an estimator for the integrated covariance of X:

$$(\widehat{X}_t)_{t\in\Lambda_2} \longrightarrow \widehat{\int_0^T q_s ds},$$

where q is defined by $q_{iji'j'} = \sum_{k,l} Q^k_{ij} Q^l_{i'j'}$.

(3) under some parametric specification of X, e.g., being an affine S_d^+-valued process, certain parameters associated to the covariance of X and the correlation between X and Y can then be determined from the previously defined estimators.

In order to perform the first step (1), that is, the non-parametric pathwise covariance estimation, we rely on (a modification of) the Fourier method introduced in [8, 9]. In order to describe its main idea, let us first introduce some notation: for an $L^1[0,T]$ function f we denote its Fourier coefficients for $k \in \mathbb{Z}$ by

$$\mathcal{F}(f)(k) = \frac{1}{T}\int_0^T f(t) e^{-\mathsf{i}\frac{2\pi}{T}kt} dt.$$

The Fourier method is now best described by the following steps. Notice that – as outlined in Section 2 – we could perform these steps for any orthonormal system in $L^2([0,T])$.

(1a) Recover from a discrete observation of Y an estimator for the Fourier coefficients of the components of the path $t \to \rho(X_t(\omega))$ for some continuous invertible function $\rho : S_d \to S_d$. In other words, find an estimator for

$$\mathcal{F}(\rho(X))(k) = \frac{1}{T}\int_0^T \rho(X_t) e^{-\mathsf{i}\frac{2\pi}{T}kt} dt = \langle e_k, \rho(X)\rangle\,.$$

(1b) Use Fourier-Féjer inversion to reconstruct the path of $t \to \rho(X_t)$. In fact, by Féjer's theorem

$$\widehat{\rho(X)}^N_t := \sum_{k=-N}^{N} \left(1 - \frac{|k|}{N}\right) \mathcal{F}(\rho(X))(k) e^{\mathsf{i}\frac{2\pi}{T}kt}$$

converges uniformly (and in L^2) to $t \mapsto \rho(X_t)$ on $[0,T]$ if $t \to X_t$ is continuous. If X has càdlàg paths, then the limit is given by $\frac{\rho(X_t)+\rho(X_{t-})}{2}$. Due to central limit theorems on the fine grid Λ_1 we make errors in the reconstruction of $\rho(X)$ of size

$$\sum_{k=-N}^{N} \left(1 - \frac{|k|}{N}\right) \mathcal{E}_k e^{\mathsf{i}\frac{2\pi}{T}kt}\,,$$

where $\mathcal{E}_k$ is a sequence of error random variables, which are approximately conditionally Gaussian with variance of order $\mathcal{O}(n^{-1})$. Hence it does not make sense to use *all* Fourier coefficients from $-\lfloor nT \rfloor$ to $\lfloor nT \rfloor$, but there will appear a subtle relationship between the sum of the errors, the size of N with respect to n and the rate of the central limit theorem for the reconstruction. See also the remarks in Section 2.

(1c) Invert the function ρ to obtain an estimator $\widehat{X}$ of the realized path of X.

For the second step (2), we can rely on existing estimators for the realized integrated covariance, into which we plug the estimator $\widehat{X}$ obtained in the first step.

In the third step (3) we focus on particular parametric specifications of X. One particularly tractable class is the class of affine models. In this case the quadratic variation of X satisfies

$$\langle X^c_{ii}, X^c_{jj} \rangle_T = 4\alpha_{ij} \int_0^T X_{s,ij} ds, \quad \alpha \in S_d^+,$$

such that knowing an estimator for $\langle X^c_{ii}, X^c_{jj} \rangle_T$ and an estimator for

$$\langle Y^c_i, Y^c_j \rangle_T = \int_0^T X_{s,ij} ds,$$

namely $\int_0^T \widehat{X}_{s,ij} ds$ allows to estimate the parameter α_{ij}.

5. Estimating Fourier coefficients

In this section we focus on step (1a), i.e., on how the Fourier coefficients of $t \mapsto \rho(X_t)$ can be estimated from discrete observations of Y. Realizing that the only difference with respect to estimators for integrated (functions of the) realized covariance are the terms $e^{-\mathrm{i}\frac{2\pi}{T}kt}$ in the integral for the Fourier coefficients, we can make use of (Fourier basis modified) jump robust estimators like

- the power variation estimators considered by Barndorff-Nielsen et al. [1],
- estimators for the realized Laplace transform of volatility introduced by Tauchen and Todorov [12] and
- other jump robust specifications, as for example considered in [5, Theorem 5.3.5]

In order to introduce those estimators, let us start by making some assumptions on the observations of the log-price process Y. Throughout let $T > 0$ be fixed and suppose that the time grids of observations for all components of Y in $[0, T]$ are equal and equidistant, i.e.,

$$t^n_m = \frac{m}{n}, \quad m = 0, \ldots, \lfloor nT \rfloor .$$

The increments of a process Z with respect to the above time grid are denoted by $\Delta^n_m Z = Z_{t^n_m} - Z_{t^n_{m-1}}$.

Remark 5.1. If grids are non-equidistant and non-equal for different coordinates it might be wise to use estimators, whose input are more continuous quantities than increments, e.g., Fourier coefficients. This is outlined for instance in [8, 9]. In any case our method will provide as a result continuous path functionals such as Fourier coefficients after the first estimation procedure.

The estimators for the Fourier coefficients that we consider are of the form

$$V(Y, g, k)^n_T = \frac{1}{n} \sum_{m=1}^{\lfloor nT \rfloor} e^{-\mathrm{i}\frac{2\pi}{T}kt^n_{m-1}} g(\sqrt{n}\Delta^n_m Y), \tag{5.1}$$

for some function $g : \mathbb{R}^d \to S_d$ and we write

$$V(Y,g)_T^{n,N} := (V(Y,g,-N)_T^n, \dots, V(Y,g,0)_T^n, \dots, V(Y,g,N)_T^n)^\top .$$

Note that the 0-th Fourier coefficient $V(Y,g,0)_T^n$ corresponds to estimators for integrated (functions of the) covariance, as the power variation estimators, considered by Barndorff-Nielsen et al. (see, e.g., [1]), but also the realized Laplace transform estimator, introduced by Tauchen and Todorov [12]. Indeed in these cases the function g is given by

$$g : \mathbb{R}^d \to S_d, \quad (x_1, \dots, x_d)^\top \mapsto (|x_i|^r |x_j|^s)_{i,j \in \{1,\dots,d\}}, \quad r+s>0$$

and

$$g : \mathbb{R}^d \to S_d, \quad (x_1, \dots, x_d)^\top \mapsto (\cos(x_i + x_j))_{i,j \in \{1,\dots,d\}}$$

respectively.

Our first aim is to study asymptotic properties of $V(Y,g)_T^{n,N}$, for which we rely to a large extent on the results of [1], [4] and [5]. The following assumptions on the function g, needed to establish consistency and a central limit theorem, are also taken from [1]:

Assumption (J). *The function g is continuous with at most polynomial growth.*

Assumption (K). *The function g is even and continuously differentiable with partial derivatives having at most polynomial growth.*

Assumption (K'). *The function g is even, with at most polynomial growth and C^1 outside a subset B of $\mathbb{R}^d$ which is a finite union of affine hyperplanes. With $d(x,B)$ denoting the distance between $x \in \mathbb{R}^d$ and B, we have for some $w \in (0,1]$ and $p \geq 0$*

$$\begin{aligned} &x \in B^c \Rightarrow \|\nabla g(x)\| \leq C(1+\|x\|^p)\left(1 + \frac{1}{d(x,B)^{1-w}}\right) \\ &x \in B^c,\ \|y\| \leq (1 \wedge \frac{d(x,B)}{2}) \\ &\qquad \Rightarrow \|\nabla f(x+y) - \nabla f(x)\| \leq C\|y\|(1+\|x\|^p + \|y\|^p)\left(1+\frac{1}{d(x,B)^{2-w}}\right). \end{aligned}$$

Remark 5.2. The conditions of Assumption (K') are especially designed to accommodate the functions

$$(x_1, \dots, x_d)^\top \mapsto (|x_i|^r |x_j|^s)_{i,j \in \{1,\dots,d\}}$$

for $r+s<1$, which correspond to the jump robust power variation estimators.

In the case when Y is a pure diffusion process the results of [1] carry directly over to the Fourier basis modified statistics $V(Y,g,k)_T^n$. In the case of jumps, the respective assertions of [5, Theorem 3.4.1, Theorem 5.3.5 and Theorem 5.3.6] can also be directly transfered to $V(Y,g,k)_T^n$. A sufficient condition which allows to incorporate jumps and which is also satisfied by the assumptions of the cited theorems, relates the function g with the jump activity of (a localized version of) Y and is stated in Assumption $(L(\eta))$ below. Let us denote by D^Y the diffusion part of Y with respect to some truncation function χ', i.e.,

$$D_t^Y(\chi') = y + \int_0^t \left(b_s^Y + \int (\chi'(\xi) - \chi(\xi)) K_s(d\xi) \right) ds + \int_0^t \sqrt{X_{s-}} dZ_s. \tag{5.2}$$

Then we shall require that the L^1-norm of $g(\sqrt{n}\Delta_m^n Y(p)) - g(\sqrt{n}\Delta_m^n D^{Y(p)}(\chi'))$ goes sufficiently fast to 0 uniformly in m, where $Y(p)$ denotes a localized version of Y.

Assumption $(L(\eta))$. *Let χ' be a truncation function such that the modified drift of Y*

$$b^Y(\chi') = b^Y + \int (\chi'(\xi) - \chi(\xi))K(d\xi) \tag{5.3}$$

is càglàd. Moreover, suppose that there exists an increasing sequence of stopping times (τ_p) with $\lim_p \tau_p = \infty$ a.s. and processes $Y(p)$ such that for $t < \tau_p$

$$Y(p)_t = Y_t \text{ a.s and } D_t^Y(\chi') = D_t^{Y(p)}(\chi') \text{ a.s.} \tag{5.4}$$

For $\eta \geq 0$, we then have for all p

$$\lim_{n\to\infty} \sup_{1\leq m\leq \lfloor nT \rfloor} n^\eta \mathbb{E}\left[\left\| g(\sqrt{n}\Delta_m^n Y(p)) - g(\sqrt{n}\Delta_m^n D^{Y(p)}(\chi'))\right\|\right] = 0.$$

In Section 5.2, we shall give precise examples of g for which this condition is satisfied.

In the following section and throughout the paper, C always denotes some constant which can vary from line to line.

5.1. Consistency and a central limit theorem for estimators of the Fourier coefficients.

The following result for the statistics $V(Y,g)_T^{n,N}$ is a slight modification of [1, Theorem 2.1. and Theorem 2.3] when Y is a diffusion and of [5, Theorem 3.4.1, Theorem 5.3.5 and Theorem 5.3.6] in the general case.

For its formulation we need some further notation: let $f : [0,T] \to S_d$ be some $L^1([0,T])$ function. Then we denote the $(2N+1)d \times d$ dimensional "vector" of Fourier coefficients (the "coordinates" of Section 2) by

$$\mathcal{F}^N(f) = (\mathcal{F}(f)(-N), \ldots, \mathcal{F}(f)(0), \ldots, \mathcal{F}(f)(N))^\top .$$

Moreover, for a function $h : \mathbb{R}^d \to \mathbb{R}^m$ and a d-dimensional normally distributed random variable U with mean 0 and covariance X, the first moment of $h(U)$ is denoted by $\rho_h(X)$, i.e.,

$$\rho_h(X) = \mathbb{E}[h(U)], \quad U \sim \mathcal{N}(0, X).$$

By $\mathcal{F}(\rho_h(X))(k)$ we then mean

$$\mathcal{F}(\rho_h(X))(k) = \frac{1}{T}\int_0^T \rho_h(X_t)e^{-\mathrm{i}\frac{2\pi}{T}kt}dt.$$

Theorem 5.3. *Under Assumptions (H), (J) and $(L(0))$, we have*

$$V(Y,g)_T^{n,N} \xrightarrow{\mathbb{P}} T\mathcal{F}^N(\rho_g(X)).$$

Moreover, under the assumption $(H1)$ and (K) or (K') and $(L(\frac{1}{2}))$, the $\mathbb{C}^{(2N+1)d\times d}$-valued random variable

$$\sqrt{n}\left(V(Y,g)_T^{n,N} - T\mathcal{F}^N(\rho_g(X))\right)$$

converges for $n \to \infty$ stably in law to an $\mathcal{F}$-conditional Gaussian random variable defined on an extension of the original probability space with mean 0 and covariance

$$C^{kk'}_{iji'j'} := \int_0^T \left(\rho_{g_{ij}g_{i'j'}}(X_s) - \rho_{g_{ij}}(X_s)\rho_{g_{i'j'}}(X_s)\right) e^{-\mathrm{i}\frac{2\pi}{T}(k-k')s} ds,$$

where $i, j, i', j' \in \{1, \dots, d\}$ and $k, k' \in \{-N, \dots, N\}$.

Remark 5.4.

- (i) Stable convergence in law for a sequence of random variables (U_n) to a limit U (defined on an extension of $(\Omega, \mathcal{F}, \mathbb{P})$) means that, for any bounded continuous function f and any bounded $\mathcal{F}$-measurable random variable V, we have

  $$\lim_{n\to\infty} \mathbb{E}\left[V f(U_n)\right] = \mathbb{E}\left[V f(U)\right].$$

- (ii) The above convergence results do not only hold for T fixed, but we have

  $$V(Y, g)^{n,N}_T \xrightarrow{\mathbb{P}} T\mathcal{F}^N(\rho_g(X))$$

  locally uniformly in T and also stable convergence process-wise.[1] The latter means that

  $$\sqrt{n}\left(V(Y, g)^{n,N}_T - T\mathcal{F}^N(\rho_g(X))\right)$$

  converges stably in law to a process $U(g, N)$ given componentwise by

  $$U(g, N)^k_{ij,T} = \sum_{k'=1}^{2(2N+1)} \sum_{i',j'=1}^{d} \int_0^T \delta^{kk'}_{s,ij,i'j'} dW^{k'}_{s,i'j}, \tag{5.5}$$

  where

  $$\sum_{r=1}^{2(2N+1)} \sum_{p,q=1}^{d} \int_0^T \delta^{kr}_{s,ij,pq} \overline{\delta^{k'r}_{s,i'j',pq}} = C^{kk'}_{iji'j'}.$$

  Here, W is a $2(2N + 1)d \times d$-dimensional Brownian motion which is defined on an extension of the probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$ and is independent of the σ-field $\mathcal{F}$.
- (iii) The above theorem has been proved in [1] in a pure diffusion setting and $k = 0$. Inclusion of jumps has been considered (in the one-dimensional case) in [2] and [13] for $g = |x|^r$ and in [12] for $g = \cos(x)$. More general functions (also for the case $k = 0$) are treated in [4] and [5, Theorem 3.4.1, Theorem 5.3.5 and Theorem 5.3.6].
- (iv) In the examples $g(x) = |x|^r$ and $g(x) = \cos(x)$, the function $\rho_g(x)$ corresponds to

  $$\rho_{(x\mapsto|x|^r)}(x) = |x|^{\frac{r}{2}}\mathbb{E}\left[|U|^r\right], \; U \sim \mathcal{N}(0, 1)$$

  and

  $$\rho_{(x\mapsto\cos(x))}(x) = e^{-\frac{1}{2}x},$$

  respectively.

---

[1] Here, $\mathcal{F}(f)(k)$ is defined for variable T.

Proof. We consider the one-dimensional case, i.e., $g : \mathbb{R} \to \mathbb{R}$, since the multi-dimensional case follows from it immediately in a methodological sense. Moreover, by the so called *localization procedure* as described in [1, Section 3] or [5, Section 4.4.1], we can strengthen assumption (H) on Y together with Condition (5.3) to (SH) (see [1, Section 3]), that is, $b^Y(\chi')$ defined in (5.3), and X are supposed to be uniformly bounded by a constant. Similarly, assumption $(H1)$ on X is strengthened to $(SH1)$, that is, (SH) holds and the differential characteristics of the drift and the diffusion part of $\sqrt{X}$ and (3.2), that is,

$$\left(\int_{S_d} (\sqrt{X_{t-} + \xi} - \sqrt{X_{t-}})\, F_t(d\xi)\right)_{t \geq 0}$$

are bounded by a constant (compare [1, Assumption $(SH1)$] and [5, Assumption 4.4.7 (or (SK))]).

Let us introduce the following notation

$$\begin{aligned}
\mathcal{B}^N(t) &:= (e^{-\mathrm{i}\frac{2\pi}{T}(-N)t}, \ldots, 1, \ldots, e^{-\mathrm{i}\frac{2\pi}{T}(N)t})^\top, \\
\beta^n_m &:= \sqrt{n}\sqrt{X_{t^n_{m-1}}}\Delta^n_m Z, \\
\rho^n_m(g) &:= \rho_g(X_{t^n_m})
\end{aligned}$$

and let us split

$$\begin{aligned}
&\sqrt{n}\left(V(Y,g)^{n,N}_T - T\mathcal{F}^N(\rho_g(X))\right) \\
&\quad = \sum_{m=1}^{\lfloor nT \rfloor} \underbrace{\left(\frac{1}{\sqrt{n}}\mathcal{B}^N(t^n_{m-1})g(\sqrt{n}\Delta^n_m Y) - \sqrt{n}\int_{t^n_{m-1}}^{t^n_m} \mathcal{B}^N(s)\rho_g(X_s)ds\right)}_{\eta^n_m}
\end{aligned}$$

into

$$\sum_{m=1}^{\lfloor nT \rfloor} (\eta^{n,1}_m + \eta^{n,2}_m + \eta^{n,3}_m)\,,$$

where

$$\begin{aligned}
\eta^{n,1}_m &= \frac{1}{\sqrt{n}}\mathcal{B}^N(t^n_{m-1})(g(\beta^n_m) - \rho^n_{m-1}(g))\,, \\
\eta^{n,2}_m &= \sqrt{n}\int_{t^n_{m-1}}^{t^n_m} (\mathcal{B}^N(t^n_{m-1})\rho^n_{m-1}(g) - \mathcal{B}^N(s)\rho_g(X_s))ds\,, \\
\eta^{n,3}_m &= \frac{1}{\sqrt{n}}\mathcal{B}^N(t^n_{m-1})(g(\sqrt{n}\Delta^n_m Y) - g(\beta^n_m))\,.
\end{aligned}$$

We divide the proof into several steps: the *first step*, which is subject of Proposition 5.5 below, consists in dealing with

$$U^n_T(g,N) := \sum_{m=1}^{\lfloor nT \rfloor} \eta^{n,1}_m = \frac{1}{\sqrt{n}}\sum_{m=1}^{\lfloor nT \rfloor} \mathcal{B}^N(t^n_{m-1})(g(\beta^n_m) - \rho^n_{m-1}(g)). \tag{5.6}$$

As proved in Proposition 5.5 below, $(U^n(g,N))_{n\in\mathbb{N}}$ converges stably in law to the process $U(g,N)$, defined in (5.5).

For the central limit theorem the *second step* consists in showing that

$$\sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,2} = \sum_{m=1}^{\lfloor nT \rfloor} \sqrt{n} \int_{t_{m-1}^n}^{t_m^n} (\mathcal{B}^N(t_{m-1}^n)\rho_{m-1}^n(g) - \mathcal{B}^N(s)\rho_g(X_s))ds$$

converges in probability to 0 uniformly in T. This can be shown similarly as in [5, Section 5.3.3, B]. Note that for the consistency result it is enough that

$$\sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{\sqrt{n}} \eta_m^{n,2} = \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} \mathcal{B}^N(t_{m-1}^n)\rho_{m-1}^n(g) - \int_0^T \mathcal{B}^N(s)\rho_g(X_s)ds.$$

converges in probability to 0 uniformly in T, which is implied by Riemann integrability.

In the *third step* we finally consider

$$\sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,3} = \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{\sqrt{n}} \mathcal{B}^N(t_{m-1}^n)(g(\sqrt{n}\Delta_m^n Y) - g(\beta_m^n)),$$

which we split into $\sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,31} + \eta_m^{n,32}$, with

$$\begin{aligned} \eta_m^{n,31} &= \frac{1}{\sqrt{n}} \mathcal{B}^N(t_{m-1}^n)(g(\sqrt{n}\Delta_m^n D^Y(\chi')) - g(\beta_m^n)), \\ \eta_m^{n,32} &= \frac{1}{\sqrt{n}} \mathcal{B}^N(t_{m-1}^n)(g(\sqrt{n}\Delta_m^n Y) - g(\sqrt{n}\Delta_m^n D^Y(\chi'))), \end{aligned}$$

where $D^Y(\chi')$ denotes the diffusion part of Y defined in (5.2). For the central limit theorem to hold true, $\sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,31}$ has to converge to 0 in probability uniformly in T. To this end, it suffices to prove

$$\sum_{m=1}^{\lfloor nT \rfloor} \mathbb{E}\left[\|\eta_m^{n,31}\|^2\right] \to 0\,, \tag{5.7}$$

$$\sum_{m=1}^{\lfloor nT \rfloor} \mathbb{E}\left[\eta_m^{n,31} | \mathcal{F}_{t_{m-1}^n}\right] \xrightarrow{\mathbb{P}} 0 \tag{5.8}$$

holds true uniformly in T. Indeed, (5.7) implies (compare [5, Lemma 2.2.11])

$$\sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,31} - \mathbb{E}\left[\eta_m^{n,31} | \mathcal{F}_{t_{m-1}^n}\right] \xrightarrow{\mathbb{P}} 0$$

and thus together with (5.8), we have $\sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,31} \xrightarrow{\mathbb{P}} 0$ (uniformly in T). Both requirements (5.7) and (5.8) are met under the conditions on X, Y and g and the proof of (5.7) can be found in [1, Proof of Theorem 5.1] and (5.8) is shown in [5, Section 5.3.3, C]. For the consistency result only (5.7) has to be satisfied, which holds under the less restrictive assumptions (H) and (J) (see [1, Theorem 5.1]). Indeed, we only have to show that

$$\sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{\sqrt{n}} \eta_m^{n,31} \xrightarrow{\mathbb{P}} 0,$$

which follows from the Cauchy-Schwarz inequality and (5.7), since

$$\sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{\sqrt{n}} \mathbb{E}\left[\|\eta_m^{n,31}\|\right] \leq \left(T \sum_{m=1}^{\lfloor nT \rfloor} \mathbb{E}\left[\|\eta_m^{n,31}\|^2\right] \right)^{\frac{1}{2}} \to 0.$$

Finally, according to the assumptions $(L(0))$ or $(L(\frac{1}{2}))$, respectively, we have for all $\varepsilon > 0$

$$\mathbb{P}\left[\left\| \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{\sqrt{n}} \eta_m^{n,32} \right\| > \varepsilon \right] \leq \mathbb{P}\left[\tau_p \leq T\right] + \mathbb{P}\left[\left\| \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{\sqrt{n}} \eta_m^{n,32}(p) \right\| > \varepsilon \right], \tag{5.9}$$

and

$$\mathbb{P}\left[\left\| \sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,32} \right\| > \varepsilon \right] \leq \mathbb{P}\left[\tau_p \leq T\right] + \mathbb{P}\left[\left\| \sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,32}(p) \right\| > \varepsilon \right], \tag{5.10}$$

where

$$\eta_m^{n,32}(p) = \mathcal{B}^N(t_{m-1}^n)(g(\sqrt{n}\Delta_m^n Y(p)) - g(\sqrt{n}\Delta_m^n D^{Y(p)}(\chi'))).$$

and $Y(p)$ is defined in (5.4). Due to $(L(0))$ or $(L(\frac{1}{2}))$, respectively, the second term on the right hand side of (5.9) and (5.10) respectively tends to 0 as $n \to \infty$ for all p. Since $\mathbb{P}\left[\tau_p \leq T\right] \to 0$ as $p \to \infty$, we deduce

$$\mathbb{P}\left[\left\| \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{\sqrt{n}} \eta_m^{n,32} \right\| > \varepsilon \right] \to 0$$

and

$$\mathbb{P}\left[\left\| \sum_{m=1}^{\lfloor nT \rfloor} \eta_m^{n,32} \right\| > \varepsilon \right] \to 0$$

respectively, which completes the proof. □

The following proposition is an application of [1, Proposition 4.1] or [5, Theorem 4.2.1], whose proof we state for convenience. Again we here only consider the one-dimensional case, i.e., Y and X are one-dimensional and $g : \mathbb{R} \to \mathbb{R}$.

Proposition 5.5. *Assume that the process X is càdlàg and bounded by a constant. Let g be a function of at most polynomial growth. Then the sequence of processes $(U_T^n(g,N))_{n\in\mathbb{N}}$ defined in (5.6) is C-tight. Moreover, if g is even, then it converges stably in law to the process $U(g,N)$ given componentwise for $k \in \{-N,\dots,N\}$ by*

$$U(g,N)_T^k = \sum_{k'=1}^{2(2N+1)} \int_0^T \delta_s^{kk'} dW_s^{k'}, \tag{5.11}$$

where

$$\sum_{r=1}^{2(2N+1)} \delta_s^{kr} \overline{\delta_s^{k'r}} = (\rho_{g^2}(X_s) - (\rho_g(X_s))^2) e^{-\mathrm{i}\frac{2\pi}{T}(k-k')s}$$

and W is a $2(2N+1)$-dimensional Brownian motion which is defined on an extension of the probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$ and is independent of the σ-field $\mathcal{F}$.

Proof. Let us write

$$U^n_T(g,N) = \sum_{m=1}^{\lfloor nT \rfloor} \zeta^n_m$$

with

$$\zeta^n_m := \eta^{n,1}_m = \frac{1}{\sqrt{n}} \mathcal{B}^N(t^n_{m-1})(g(\beta^n_m) - \rho^n_{m-1}(g)).$$

In particular, for $k \in \{-N, \dots, N\}$, let

$$\begin{aligned} \zeta^n_m(k) &= \frac{1}{\sqrt{n}} e^{-\mathrm{i}\frac{2\pi}{T} k t^n_{m-1}} (g(\beta^n_m) - \rho^n_{m-1}(g)), \\ U^n_T(g,N)(k) &= \sum_{m=1}^{\lfloor nT \rfloor} \zeta^n_m(k). \end{aligned}$$

Observe that

$$\mathbb{E}\left[\zeta^n_m | \mathcal{F}_{t^n_{m-1}}\right] = 0 \tag{5.12}$$

and

$$\mathbb{E}\left[(g(\beta^n_m) - \rho^n_{m-1}(g))^2 | \mathcal{F}_{t^n_{m-1}}\right] = \rho^n_{m-1}(g^2) - (\rho^n_{m-1}(g))^2.$$

Hence we have, for $k, l \in \{-N, \dots, N\}$,

$$\begin{aligned} &\mathbb{E}\left[\mathrm{Re}\, \zeta^n_m(k) \mathrm{Re}\, \zeta^n_m(l) |\, \mathcal{F}_{t^n_{m-1}}\right] \\ &\quad = \frac{1}{n} \cos\left(-\frac{2\pi}{T} k t^n_{m-1}\right) \cos\left(-\frac{2\pi}{T} l t^n_{m-1}\right) (\rho^n_{m-1}(g^2) - (\rho^n_{m-1}(g))^2), \\ &\mathbb{E}\left[\mathrm{Im}\, \zeta^n_m(k) \mathrm{Im}\, \zeta^n_m(l) |\, \mathcal{F}_{t^n_{m-1}}\right] \\ &\quad = \frac{1}{n} \sin\left(-\frac{2\pi}{T} k t^n_{m-1}\right) \sin\left(-\frac{2\pi}{T} l t^n_{m-1}\right) (\rho^n_{m-1}(g^2) - (\rho^n_{m-1}(g))^2), \\ &\mathbb{E}\left[\mathrm{Im}\, \zeta^n_m(k) \mathrm{Re}\, \zeta^n_m(l) |\, \mathcal{F}_{t^n_{m-1}}\right] \\ &\quad = \frac{1}{n} \sin\left(-\frac{2\pi}{T} k t^n_{m-1}\right) \cos\left(-\frac{2\pi}{T} l t^n_{m-1}\right) (\rho^n_{m-1}(g^2) - (\rho^n_{m-1}(g))^2). \end{aligned}$$

Thus by Riemann integrability and the càdlàg assumption on X, it follows that

$$
\begin{aligned}
&\sum_{m=0}^{\lfloor nT \rfloor} \mathbb{E}\left[\operatorname{Re}\zeta_m^n(k)\operatorname{Re}\zeta_m^n(l)|\mathcal{F}_{t_{m-1}^n}\right] \\
&\qquad \to \int_0^T \cos\left(-\frac{2\pi}{T}ks\right)\cos\left(-\frac{2\pi}{T}ls\right)(\rho_{g^2}(X_s)-(\rho_g(X_s))^2)ds\,, \\
&\sum_{m=0}^{\lfloor nT \rfloor} \mathbb{E}\left[\operatorname{Im}\zeta_m^n(k)\operatorname{Im}\zeta_m^n(l)|\mathcal{F}_{t_{m-1}^n}\right] \\
&\qquad \to \int_0^T \sin\left(-\frac{2\pi}{T}ks\right)\sin\left(-\frac{2\pi}{T}ls\right)(\rho_{g^2}(X_s)-(\rho_g(X_s))^2)ds\,, \\
&\sum_{m=0}^{\lfloor nT \rfloor} \mathbb{E}\left[\operatorname{Im}\zeta_m^n(k)\operatorname{Re}\zeta_m^n(l)|\mathcal{F}_{t_{m-1}^n}\right] \\
&\qquad \to \int_0^T \sin\left(-\frac{2\pi}{T}ks\right)\cos\left(-\frac{2\pi}{T}ls\right)(\rho_{g^2}(X_s)-(\rho_g(X_s))^2)ds\,,
\end{aligned} \tag{5.13}
$$

as $n \to \infty$. Due to the polynomial growth of g we further obtain

$$
\mathbb{E}\left[\|\zeta_m^n\|^4|\mathcal{F}_{t_{m-1}^n}\right] \leq \frac{C}{n^2}, \tag{5.14}
$$

which already implies tightness. If g is even, we have due to $\Delta_m^n Z = -\Delta_m^n Z$

$$
\mathbb{E}\left[\zeta_m^n \Delta_m^n Z|\mathcal{F}_{t_{m-1}^n}\right] = 0. \tag{5.15}
$$

Finally let N be any bounded martingale which is orthogonal to Z. Then we obtain by the same arguments is in [1, Proposition 4.1] that

$$
\mathbb{E}\left[\zeta_m^n \Delta_m^n N|\mathcal{F}_{t_{m-1}^n}\right] = 0. \tag{5.16}
$$

Due to (5.12), (5.13), (5.14), (5.15) and (5.16), [6, Theorem IX.7.28] (see also [5, Theorem 2.2.15]) now implies that for $k, l \in \{-N, \ldots, N\}$

$$
\begin{aligned}
\operatorname{Re} U_T^n(g,N)(k) &\overset{\mathcal{L}-s}{\to} \sum_{j=1}^{2(2N+1)} \int_0^T w_{s,kj}^{\mathrm{Re}} dW_{s,j}, \\
\operatorname{Im} U_T^n(g,N)(k) &\overset{\mathcal{L}-s}{\to} \sum_{j=1}^{2(2N+1)} \int_0^T w_{s,kj}^{\mathrm{Im}} dW_{s,j},
\end{aligned}
$$

where $\mathcal{L} - s$ means stable converges in law (see Remark 5.4 (i)) and

$$
\begin{aligned}
\sum_{j=1}^{2(2N+1)} w_{s,kj}^{\mathrm{Re}} w_{s,lj}^{\mathrm{Re}} &= \cos\left(-\frac{2\pi}{T}ks\right)\cos\left(-\frac{2\pi}{T}ls\right)(\rho_{g^2}(X_s)-(\rho_g(X_s))^2), \\
\sum_{j=1}^{2(2N+1)} w_{s,kj}^{\mathrm{Im}} w_{s,lj}^{\mathrm{Im}} &= \sin\left(-\frac{2\pi}{T}ks\right)\sin\left(-\frac{2\pi}{T}ls\right)(\rho_{g^2}(X_s)-(\rho_g(X_s))^2), \\
\sum_{j=1}^{2(2N+1)} w_{s,kj}^{\mathrm{Im}} w_{s,lj}^{\mathrm{Re}} &= \sin\left(-\frac{2\pi}{T}ks\right)\cos\left(-\frac{2\pi}{T}ls\right)(\rho_{g^2}(X_s)-(\rho_g(X_s))^2),
\end{aligned}
$$

and W is $2(2N+1)$-dimensional Brownian motion defined on an extension of the probability space $(\Omega,\mathcal{F},(\mathcal{F}_t)_{t\geq 0},\mathbb{P})$ and independent of the σ-field $\mathcal{F}$. Defining $\delta_s^{kj} = w^{\mathrm{Re}}_{s,kj} + \mathrm{i} w^{\mathrm{Im}}_{s,kj}$ yields the desired assertion. □

5.2. **Jump robust estimators.** The aim of this section is to specify classes of functions g and conditions on the jumps such that condition $(L(0))$ or $(L(\frac{1}{2}))$, respectively, is satisfied and such that the estimator $V(Y,g,k)^n_T$ given in (5.1) is robust to jumps. The conditions on g are in line with the respective assumptions in [5, Theorem 3.4.1 (a), Theorem 5.3.5 (γ) and Theorem 5.3.6, Equation 5.3.11].

Proposition 5.6. (i) *Let g be continuous with $g(x) = o(\|x^2\|)$ as $\|x\| \to \infty$. Moreover, suppose that*

$$\sup_{\omega} \int_0^T \left(\|b_t^Y\| + \|X_t\| + \int (\|\xi\|^2 \wedge 1) K_t(d\xi) \right) ds < \infty \tag{5.17}$$

and $\sup_{\omega, t\in[0,T]} \|\Delta X_t(\omega)\| < \infty$. Then for all $m \in \{1,\dots,\lfloor nt \rfloor\}$

$$\lim_{n\to\infty} \mathbb{E}\left[\left\| g(\sqrt{n}\Delta^n_m Y) - g(\sqrt{n}\Delta^n_m D^Y(\mathrm{id})) \right\| \right] = 0. \tag{5.18}$$

(ii) *Suppose that g satisfies for some $q \geq 0$ and some $0 < r \leq r' < 1$*

$$\|g(x) - g(y)\| \leq C(1+\|y\|^q)(\|x-y\|^r + \|x-y\|^{r'}). \tag{5.19}$$

Moreover, let $\beta \in [0,2]$ and assume that for all $t \in [0,T]$

$$\mathbb{E}\left[\int_{\|\xi\|\leq 1} \|\xi\|^\beta K_t(d\xi) \right] < \infty, \tag{5.20}$$

$$\mathbb{E}\left[\int_{\|\xi\|> 1} \|\xi\| K_t(d\xi) \right] < \infty \tag{5.21}$$

and that $b^Y(\chi 1_{\{\beta>1\}}) = b^Y + \int (\chi(\xi)1_{\{\beta>1\}} - \chi(\xi)) K(d\xi)$ as defined in (5.3) and X is uniformly bounded.

(a) If $\beta \leq r < 1$, then there exists some $\kappa \in (1, \frac{1}{r'}]$ such that

$$\mathbb{E}\left[\left\| g(\sqrt{n}\Delta^n_m Y) - g\left(\sqrt{n}\Delta^n_m D^Y(0)\right) \right\| \right] \leq C \frac{1}{n^{\frac{1}{\kappa} - \frac{r'}{2}}}, \tag{5.22}$$

for all $m \in \{1,\dots,\lfloor nt \rfloor\}$.

(b) If $\beta > r$, then

$$\mathbb{E}\left[\left\| g(\sqrt{n}\Delta^n_m Y) - g\left(\sqrt{n}\Delta^n_m D^Y(\chi 1_{\{\beta>1\}})\right) \right\| \right] \leq C \frac{1}{n^{r\frac{2-\beta}{2\beta}}},$$

for all $m \in \{1,\dots,\lfloor nt \rfloor\}$.

Remark 5.7. (i) Another sufficient condition to obtain (5.18) is

$$\mathbb{E}\left[(b_t^Y)^2 + \int \xi^2 K_t(d\xi) + X_t \right] < \infty \tag{5.23}$$

for all $t \in [0,T]$. Note also that we can localize Y and consider processes $(Y(p))_p$ for which (5.17) or (5.23) is automatically satisfied (see e.g. [5, Lemma 3.4.5]). Assertion (i) then holds true for $Y(p)$, which implies that $(L(0))$ is satisfied without further conditions on the characteristics of Y. Similarly the boundedness assumption on $b^Y(\chi 1_{\{\beta>1\}})$ and X in statement (ii) can also be obtained by localizing the original process.

(ii) Condition (5.19) is satisfied by many classes of functions, e.g., globally Hölder continuous functions or C^1 functions which satisfy $\|g(x)\| \leq C(1+\|x\|^{r'})$ for $r' \in [0,1)$ and $\|\nabla g(x)\| \leq C(1+\|x\|^q)$ for some $q \geq 0$ (compare also [5, Remark 5.3.7]). Let us also remark that, if (5.19) is satisfied with $q=0$, κ in (5.22) can be chosen to be 1.

Proof. The first assertion is proved in [5, Lemma 3.4.6]. Concerning (ii), we shall distinguish the cases $\beta \leq 1$ and $\beta > 1$ and set without loss of generality $\chi(\xi) = 1_{\{\|\xi\|\leq 1\}}\xi$. Due to the assumption on g, we have

$$\begin{aligned}
&\mathbb{E}\left[\left\|g(\sqrt{n}\Delta_m^n Y) - g\left(\sqrt{n}\Delta_m^n D^Y(\chi 1_{\{\beta>1\}})\right)\right\|\right] \\
&\leq C\mathbb{E}\Bigg[(1+\|\sqrt{n}\Delta_m^n D^Y(\chi 1_{\{\beta>1\}})\|^q)\times \\
&\qquad \times\Bigg(\left\|\sqrt{n}\int_{t_{m-1}^n}^{t_m^n}\int_{\{\|\xi\|\leq 1\}}\xi(\mu^Y(d\xi,ds)-1_{\{\beta>1\}}K_s(d\xi)ds)\right\|^r + \\
&\qquad + \left\|\sqrt{n}\int_{t_{m-1}^n}^{t_m^n}\int_{\{\|\xi\|>1\}}\xi\mu^Y(d\xi,ds)\right\|^r + \\
&\qquad + \left\|\sqrt{n}\int_{t_{m-1}^n}^{t_m^n}\int_{\{\|\xi\|\leq 1\}}\xi(\mu^Y(d\xi,ds)-1_{\{\beta>1\}}K_s(d\xi)ds)\right\|^{r'} + \\
&\qquad + \left\|\sqrt{n}\int_{t_{m-1}^n}^{t_m^n}\int_{\{\|\xi\|>1\}}\xi\mu^Y(d\xi,ds)\right\|^{r'}\Bigg)\Bigg].
\end{aligned} \tag{5.24}$$

Applying Hölder's inequality, we get for $1 < \iota, \kappa < \infty$ such that $\frac{1}{\iota}+\frac{1}{\kappa}=1$

$$\begin{aligned}
&\mathbb{E}\left[\|\sqrt{n}\Delta_m^n D^Y(\chi 1_{\{\beta>1\}})\|^q J^k\right] \\
&\quad \leq \mathbb{E}\left[\|\sqrt{n}\Delta_m^n D^Y(\chi 1_{\{\beta>1\}})\|^{q\iota}\right]^{\frac{1}{\iota}} \mathbb{E}\left[J^{k\kappa}\right]^{\frac{1}{\kappa}}.
\end{aligned}$$

Here, J stands for each of the above expressions of the jumps and k corresponds to r or r'. Due to our assumptions on $b^Y(\chi 1_{\{\beta>1\}})$ and X and as a consequence of Hölder's, Jensen's and Burkholder-Davis-Gundy's inequality the first expectation is bounded by a constant $C_{q,\iota}$ for all $q>0$ and $1<\iota<\infty$.

Let us now consider for some $\alpha \in (0,1]$

$$\mathbb{E}\left[\left\|\int_{t_{m-1}^n}^{t_m^n}\int_{\{\|\xi\|>1\}}\xi\mu^Y(d\xi,ds)\right\|^\alpha\right].$$

Then since, for $\alpha \in (0,1]$, $\|\sum_i x_i\|^\alpha \leq \sum_i \|x_i\|^\alpha$ and due to (5.21)

$$\begin{aligned}
\mathbb{E}\left[\left\|\int_{t_{m-1}^n}^{t_m^n}\int_{\{\|\xi\|>1\}}\xi\mu^Y(d\xi,ds)\right\|^\alpha\right] &\leq \mathbb{E}\left[\int_{t_{m-1}^n}^{t_m^n}\int_{\{\|\xi\|>1\}}\|\xi\|^\alpha\mu^Y(d\xi,ds)\right] \\
&= \mathbb{E}\left[\int_{t_{m-1}^n}^{t_m^n}\int_{\{\|\xi\|>1\}}\|\xi\|^\alpha K_s(d\xi)ds\right] \\
&\leq C\frac{1}{n}.
\end{aligned}$$

Consider now the term

$$\mathbb{E}\left[\left\|\int_{t^n_{m-1}}^{t^n_m}\int_{\{\|\xi\|\le 1\}}\xi(\mu^Y(d\xi,ds)-1_{\{\beta>1\}}K_s(d\xi)ds)\right\|^{\alpha}\right]$$

for $\beta\le 1$. Then by Hölder's inequality and (5.20), we have

$$\begin{aligned}\mathbb{E}\left[\left\|\int_{t^n_{m-1}}^{t^n_m}\int_{\{\|\xi\|\le 1\}}\xi\mu^Y(d\xi,ds)\right\|^{\alpha}\right] &= \left(\mathbb{E}\left[\left\|\sum_{t_{m-1}\le s<t_m}\Delta_s Y 1_{\{\Delta_s Y\le 1\}}\right\|^{\alpha\vee\beta}\right]\right)^{\frac{\alpha}{\beta}\wedge 1}\\ &\le \left(\mathbb{E}\left[\sum_{t_{m-1}\le s<t_m}\|\Delta_s Y\|^{\alpha\vee\beta}1_{\{\Delta_s Y\le 1\}}\right]\right)^{\frac{\alpha}{\beta}\wedge 1}\\ &= \left(\mathbb{E}\left[\int_{t^n_{m-1}}^{t^n_m}\int_{\{\|\xi\|\le 1\}}\|\xi\|^{\beta}K_s(d\xi)ds\right]\right)^{\frac{\alpha}{\beta}\wedge 1}\\ &\le C\frac{1}{n^{1\wedge\frac{\alpha}{\beta}}}.\end{aligned}$$

If $\beta>1$, we obtain a similar estimate by using Hölder and Burkholder-Davis-Gundy's inequality, the fact that $\frac{\beta}{2}\le 1$ and (5.20):

$$\begin{aligned}&\mathbb{E}\left[\left\|\int_{t^n_{m-1}}^{t^n_m}\int_{\{\|\xi\|\le 1\}}\xi(\mu^Y(d\xi,ds)-K_s(d\xi)ds)\right\|^{\alpha}\right]\\ &\le\left(\mathbb{E}\left[\left\|\int_{t^n_{m-1}}^{t^n_m}\int_{\{\|\xi\|\le 1\}}\xi(\mu^Y(d\xi,ds)-K_s(d\xi)ds)\right\|^{\beta}\right]\right)^{\frac{\alpha}{\beta}}\\ &\le\left(\mathbb{E}\left[\left(\int_{t^n_{m-1}}^{t^n_m}\int_{\{\|\xi\|\le 1\}}\|\xi\|^2\mu^Y(d\xi,ds)\right)^{\frac{\beta}{2}}\right]\right)^{\frac{\alpha}{\beta}}\\ &\le\left(\mathbb{E}\left[\left(\sum_{t_{m-1}\le s<t_m}\|\Delta_s Y\|^{\beta}1_{\{\Delta_s Y\le 1\}}\right)\right]\right)^{\frac{\alpha}{\beta}}\\ &\le\left(\mathbb{E}\left[\int_{t^n_{m-1}}^{t^n_m}\int_{\{\|\xi\|\le 1\}}\|\xi\|^{\beta}K_s(d\xi)ds\right]\right)^{\frac{\alpha}{\beta}}\\ &\le C\frac{1}{n^{\frac{\alpha}{\beta}}}=C\frac{1}{n^{\frac{\alpha}{\beta}\wedge 1}}.\end{aligned}$$

The last equality follows from the fact that for $\beta>1$, $\frac{\alpha}{\beta}<1$. Using these inequalities and setting α equal to $r\kappa$ and $r'\kappa'$, respectively, for some $1<\kappa\le\frac{1}{r}$ and $1<\kappa'\le\frac{1}{r'}$, we can estimate (5.24) by

$$\mathbb{E}\left[\left\|g(\sqrt{n}\Delta^n_m Y)-g\left(\sqrt{n}\Delta^n_m D^Y(\chi 1_{\{\beta>1\}})\right)\right\|\right]\le C\frac{1}{n^{\min((\frac{1}{\kappa}\wedge\frac{r}{\beta})-\frac{r}{2},(\frac{1}{\kappa'}\wedge\frac{r'}{\beta})-\frac{r'}{2})}}.$$

If $\beta\le r$, we can choose some $\kappa\in(1,\frac{1}{r'}]$ such that this expression is simplified to

$$\mathbb{E}\left[\left\|g(\sqrt{n}\Delta^n_m Y)-g\left(\sqrt{n}\Delta^n_m D^Y(\chi 1_{\{\beta>1\}})\right)\right\|\right]\le C\frac{1}{n^{\frac{1}{\kappa}-\frac{r'}{2}}}$$

and if $\beta > r$, we obtain

$$\mathbb{E}\left[\left\|g(\sqrt{n}\Delta_m^n Y) - g\left(\sqrt{n}\Delta_m^n D^Y(\chi 1_{\{\beta>1\}})\right)\right\|\right] \leq C\frac{1}{n^{r\frac{(2-\beta)}{2\beta}}}.$$

□

Corollary 5.8. (i) *Let g be continuous with $g(x) = o(\|x^2\|)$ as $\|x\| \to \infty$. Then under assumption (H) we have*

$$V(Y,g)_T^{n,N} \xrightarrow{\mathbb{P}} T\mathcal{F}^N(\rho_g(X)).$$

(ii) *Let g satisfy* (5.19) *with $r \in (0,1)$ and assume that* (5.21) *holds true. Moreover, let $\beta \in [0,1)$ and suppose that* (5.20) *is satisfied and that*

$$b^Y(0) = b^Y - \int \chi(\xi)K(d\xi)$$

is càglàd. Then under the assumptions $(H1)$ and (K) or (K') and $\frac{\beta}{2-\beta} < r < 1$, the $\mathbb{C}^{(2N+1)d\times d}$-valued random variable

$$\sqrt{n}\left(V(Y,g)_T^{n,N} - T\mathcal{F}^N(\rho_g(X))\right)$$

converges for $n \to \infty$ stably in law to an $\mathcal{F}$-conditional Gaussian random variable defined on an extension of the original probability space with mean 0 *and covariance*

$$C_{iji'j'}^{kk'} := \int_0^T \left(\rho_{g_{ij}g_{i'j'}}(X_s) - \rho_{g_{ij}}(X_s)\rho_{g_{i'j'}}(X_s)\right) e^{-\mathrm{i}\frac{2\pi}{T}(k-k')s}ds,$$

where $i,j \in \{1,\dots,d\}$ and $k,k' \in \{-N,\dots,N\}$.

Proof. Assertion (i) is a direct consequence of Theorem 5.3, Condition $(L(0))$, Proposition 5.6 (i) and Remark 5.7 (i).

Again, in view of Theorem 5.3, the proof of the second statement consists in verifying Condition $(L(\frac{1}{2}))$. Since $\beta < 1$, we can consider $b^Y(0) = b^Y - \int \chi(\xi)K(d\xi)$ which is – since it is a càglàd process – locally bounded. By the localization procedure (see [5, Lemma 4.4.8]) we can therefore consider processes $(Y(p))_p$, for which $b^{Y(p)}(0)$ and the diffusion characteristic X are uniformly bounded and (5.21) and (5.20) for $\beta < 1$ are satisfied, as required in Proposition 5.6 (ii).

This proposition then yields in the case $\frac{\beta}{2-\beta} < \beta \leq r$

$$\mathbb{E}\left[\left\|g(\sqrt{n}\Delta_m^n Y(p)) - g\left(\sqrt{n}\Delta_m^n D^{Y(p)}(0)\right)\right\|\right] \leq C\frac{1}{n^{\frac{1}{\kappa}-\frac{r'}{2}}}, \tag{5.25}$$

for some $\kappa \in (1, \frac{1}{r'}]$ and in the case $\frac{\beta}{2-\beta} < r < \beta$

$$\mathbb{E}\left[\left\|g(\sqrt{n}\Delta_m^n Y(p)) - g\left(\sqrt{n}\Delta_m^n D^{Y(p)}(0)\right)\right\|\right] \leq C\frac{1}{n^{r\frac{2-\beta}{2\beta}}}.$$

By choosing $1 < \kappa < \frac{2}{1+r'} < \frac{1}{r'}$ in (5.25), $(L(\frac{1}{2}))$ is satisfied in both cases and the assertion follows. □

Remark 5.9. (i) The specifications

$$g : \mathbb{R}^d \to S_d, \quad (x_1, \dots, x_d)^\top \mapsto (|x_i x_j|^{\frac{r}{2}})_{i,j \in \{1,\dots,d\}} \tag{5.26}$$

for $r < 1$ and

$$g : \mathbb{R}^d \to S_d, \quad (x_1, \dots, x_d)^\top \mapsto (\cos(x_i + x_j))_{i,j \in \{1,\dots,d\}}$$

are covered by these conditions. In the case of (5.26), the above corollary recovers [2, Theorem 1 (iii)], which has been proved for one dimensional jump diffusions where the jumps are described by a Lévy process. For functions satisfying (5.19) a similar statement is proved in [5, Theorem 5.3.5 (γ) and Theorem 5.3.6], however, under slightly different conditions on the jump measures (in particular, supposing the Gringelions representation of Y).

(ii) Another function which satisfies for example the above requirements and for which ρ_g is invertible and easily computable is

$$g : \mathbb{R}^d \to S_d, \quad (x_1, \dots, x_d)^\top \mapsto \left(e^{-\frac{\langle x, A_{ij} x\rangle}{2}}\right)_{ij \in \{1,\dots,d\}},$$

where $A_{ij} = e_i e_i^\top + e_j e_j^\top + e_i e_j^\top + e_j e_i^\top$ and e_i denotes the canonical basis vector. The function ρ_g is given by

$$\rho_{g_{ij}}(X) = \frac{1}{\sqrt{X_{ii} + 2X_{ij} + X_{jj} + 1}}.$$

6. ASYMPTOTIC PROPERTIES OF ESTIMATORS FOR (FUNCTIONS OF) THE INSTANTANEOUS COVARIANCE PROCESS

In this section we focus on step (1b) and in the sequel on step (1c), that is, we are interested in establishing consistency and a central limit theorem for an estimator of $\rho_g(X_t)$ and X_t respectively. The estimator for $\rho_g(X_t)$ is defined via Fourier-Féjer inversion using the above estimators for the Fourier coefficients:

$$\widehat{\rho_g(X)}_t^{n,N} = \frac{1}{T} \sum_{k=-N}^{N} \left(1 - \frac{|k|}{N}\right) e^{\mathrm{i} \frac{2\pi}{T} kt} V(Y, g, k)_T^n. \tag{6.1}$$

Once we have obtained a consistency and a central limit theorem for this estimator, we can translate these results to an estimator for X_t, which we define via

$$\widehat{X}_t^{n,N} := \rho_g^{-1}\left(\widehat{\rho_g(X)}_t^{n,N}\right) \tag{6.2}$$

provided that $\rho_g(x) : S_d \times S_d$, $x \mapsto \rho_g(x)$ is invertible.

Let us start with the following lemma which deals with convergence of Riemann sums for the Féjer kernel and which is crucial for studying asymptotic properties of the above estimators.

Lemma 6.1. *Let $n, N \in \mathbb{N}$ and denote*

$$x_0^n := -\pi < x_1^n := -\pi + \frac{1}{n} < \cdots < x_m^n := -\pi + \frac{m}{n} < x_{\lfloor n2\pi \rfloor}^n = -\pi + \frac{\lfloor n2\pi \rfloor}{n}$$

and

$$F_N(x) := \frac{1}{N+1}\frac{\sin\left((N+1)\frac{x}{2}\right)^2}{\sin\left(\frac{x}{2}\right)^2} = \sum_{k=-N}^{k=N}\left(1-\frac{|k|}{N}\right)e^{\mathrm{i}kx}. \tag{6.3}$$

Moreover, let $\gamma \in \mathbb{R}$ *and suppose that* $\lim \frac{n}{N^\gamma} = K$ *for some constant* $K > 0$ *and let* h *be a* δ*-Hölder continuous function. Then, we have*

$$\lim_{n,N\to\infty}\sum_{m=1}^{\lfloor n2\pi\rfloor}\frac{1}{n}F_N(x^n_{m-1}) = \lim_{N\to\infty}\int_{-\pi}^{\pi}F_N(x)dx = 2\pi, \tag{6.4}$$

$$\lim_{n,N\to\infty}\sum_{m=1}^{\lfloor n2\pi\rfloor}\frac{1}{n}\frac{F_N^2(x^n_{m-1})}{N} = \lim_{N\to\infty}\int_{-\pi}^{\pi}\frac{F_N^2(x)}{N}dx = \frac{4\pi}{3}, \tag{6.5}$$

$$\lim_{n,N\to\infty}\sum_{m=1}^{\lfloor n2\pi\rfloor}\frac{1}{n}\frac{F_N^2(y-x^n_{m-1})}{N}h(x^n_{m-1}) = \lim_{N\to\infty}\int_{-\pi}^{\pi}\frac{F_N^2(y-x)}{N}h(x)dx = \frac{4\pi}{3}h(y), \tag{6.6}$$

if $\gamma > 1$*. Moreover, if* $\gamma > 1$*, then*

$$\left|\sum_{m=1}^{\lfloor n2\pi\rfloor}\frac{1}{n}F_N(x^n_{m-1}) - \int_{-\pi}^{\pi}F_N(x)dx\right| \leq C\frac{1}{N^{\gamma-1}}$$

and

$$\left|\sum_{m=1}^{\lfloor n2\pi\rfloor}\frac{1}{n}h(x^n_{m-1})F_N(x^n_{m-1}) - \int_{-\pi}^{\pi}h(x)F_N(x)dx\right| \leq C\max\left(\frac{1}{N^{\gamma-1}},\frac{1}{N^{\delta\gamma}}\right).$$

Remark 6.2.

(i) In the sequel we shall consider analogues of expressions (6.5) and (6.6) on the interval $[0,T]$, that is,

$$\lim_{N\to\infty}\int_0^T\frac{F_N^2(\frac{2\pi}{T}x)}{N}dx = \frac{2T}{3},$$
$$\lim_{N\to\infty}\int_0^T\frac{F_N^2(\frac{2\pi}{T}(y-x))}{N}h(x)dx = \frac{2T}{3}h(y),$$

which can be derived similarly as in the proof below.

(ii) The expression $F_N(x)^2/N$ appropriately normalized corresponds to the so-called Jackson kernel (see e.g. [7, Section 4.2]).

Proof. We have convergence of the Riemann sums in (6.4) to the corresponding integral if the grid becomes finer within the zeros of F_N. Since the distance between 2 zeros is $\frac{2\pi}{N+1}$ ($\frac{4\pi}{N+1}$ for the 2 zeros closest to the origin), this is the case if only if $\gamma > 1$. By the same argument the left hand side of (6.5) and (6.6) converges to the corresponding integrals if and only if $\gamma > 1$. The assertion then follows from the following calculation

$$\begin{aligned}
&\int_{-\pi}^{\pi} \frac{1}{N} \sum_{k,l} \left(1 - \frac{|k|}{N}\right) \left(1 - \frac{|l|}{N}\right) e^{\mathrm{i}kx} e^{\mathrm{i}lx} dx \\
&= \frac{1}{N} \sum_{k,l} \left(1 - \frac{|k|}{N}\right) \left(1 - \frac{|l|}{N}\right) \int_{-\pi}^{\pi} e^{\mathrm{i}(k+l)x} dx \\
&= 2\pi \frac{1}{N} \sum_{k} \left(1 - \frac{|k|}{N}\right)^2 \\
&= 2\pi \frac{2N^2+1}{3N^2} \to \frac{4\pi}{3}.
\end{aligned}$$

Concerning the second statement, we have due to the mean value theorem

$$\begin{aligned}
&\left| \sum_{m=1}^{\lfloor n2\pi \rfloor} \frac{1}{n} F_N(x_{m-1}^n) - \int_{-\pi}^{\pi} F_N(x) dx \right| \\
&= \left| \sum_{m=1}^{\lfloor n2\pi \rfloor} \frac{1}{n} \left(F_N(x_{m-1}^n) - F_N(y_{m-1}^n) \right) \right| \\
&\le \sum_{m=1}^{\lfloor n2\pi \rfloor} \frac{1}{n} \left| F_N(x_{m-1}^n) - F_N(y_{m-1}^n) \right| \\
&\le \sum_{k=-\frac{(N+1)}{2}}^{\frac{(N-1)}{2}} \sum_{x_m \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} \frac{1}{n} \left| F_N(x_{m-1}^n) - F_N(y_{m-1}^n) \right|
\end{aligned}$$

with some $y_{m-1}^n \in [x_{m-1}^n, x_m^n]$. Using again the mean value theorem, we can further estimate

$$\begin{aligned}
&\sum_{x_m \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} \frac{1}{n} \left| F_N(x_{m-1}^n) - F_N(y_{m-1}^n) \right| \\
&\le \sum_{\eta_{m-1}^n \in [x_{m-1}^n, y_{m-1}^n],\, x_m \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} \frac{1}{n^2} \left| F_N'(\eta_{m-1}^n) \right| \\
&\le \max_{\eta \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} |F_N'(\eta)| \frac{2\pi n}{N+1} \frac{1}{n^2}.
\end{aligned}$$

Since

$$\max_{\eta \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} |F_N'(\eta)| \frac{1}{N} \le C \max_{x \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} |F_N(x)|,$$

and since

$$\max_{x \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} |F_N(x)| \frac{1}{N} \le C \int_{\frac{2\pi k}{N+1}}^{\frac{2\pi(k+1)}{N+1}} F_N(x) dx,$$

it follows that

$$\max_{\eta \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} |F_N'(\eta)| \le C N^2 \int_{\frac{2\pi k}{N+1}}^{\frac{2\pi(k+1)}{N+1}} F_N(x) dx.$$

Hence, we have

$$\begin{aligned}
&\sum_{k=-\frac{(N+1)}{2}}^{\frac{(N-1)}{2}} \sum_{x_m \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} \frac{1}{n}\left|F_N(x^n_{m-1}) - F_N(y^n_{m-1})\right| \\
&\leq \frac{2\pi n}{N+1}\frac{1}{n^2} \sum_{k=-\frac{(N+1)}{2}}^{\frac{(N-1)}{2}} \max_{\eta \in [\frac{2\pi k}{N+1}, \frac{2\pi(k+1)}{N+1}]} |F'_N(\eta)| \\
&\leq \frac{2\pi n}{N+1}\frac{1}{n^2} \sum_{k=-\frac{(N+1)}{2}}^{\frac{(N-1)}{2}} CN^2 \int_{\frac{2\pi k}{N+1}}^{\frac{2\pi(k+1)}{N+1}} F_N(x)dx \\
&\leq C\frac{2\pi n}{N+1}\frac{1}{n^2} N^2 \underbrace{\int_{-\pi}^{\pi} F_N(x)dx}_{=2\pi} \\
&\leq C\frac{N}{n} = C\frac{N^\gamma}{n}\frac{1}{N^{\gamma-1}},
\end{aligned}$$

which yields the assertion. Concerning

$$\left|\sum_{m=1}^{\lfloor n2\pi \rfloor} \frac{1}{n} h(x^n_{m-1})F_N(x^n_{m-1}) - \int_{-\pi}^{\pi} h(x)F_N(x)dx\right|$$

we can estimate it by

$$\begin{aligned}
&\left|\sum_{m=1}^{\lfloor n2\pi \rfloor} \frac{1}{n} h(x^n_{m-1})F_N(x^n_{m-1}) - \int_{x^n_{m-1}}^{x_m} h(x^n_{m-1})F_N(x)dx\right| + \\
&\quad + \left|\sum_{m=1}^{\lfloor n2\pi \rfloor} \int_{x^n_{m-1}}^{x^n_m} (h(x^n_{m-1}) - h(x))F_N(x)dx\right| \\
&\leq \max_m |h(x^n_{m-1})| \sum_{m=1}^{\lfloor n2\pi \rfloor} \left|\frac{1}{n}F_N(x^n_{m-1}) - \int_{x^n_{m-1}}^{x_m} F_N(x)dx\right| + \\
&\quad + \frac{C}{n^\delta}\frac{N^{\gamma\delta}}{N^{\gamma\delta}} \int_{-\pi}^{\pi} F_N(x)dx \\
&\leq C\left(\frac{1}{N^{\gamma-1}} + \frac{1}{N^{\gamma\delta}}\right) \\
&\leq C\max\left(\frac{1}{N^{\gamma-1}}, \frac{1}{N^{\gamma\delta}}\right).
\end{aligned}$$

□

6.1. **Consistency.** Using the above lemma we can now proceed to establish consistency of (6.1).

Theorem 6.3. *Let* $\gamma > 1$ *and suppose that* $\lim \frac{n}{N^\gamma} = K$ *for some constant* $K > 0$. *Under the assumptions (H), (J) and (L(0)) we have for every* $t \in [0, T]$

$$\widehat{\rho_g(X)}^{n,N}_t \xrightarrow{\mathbb{P}} \frac{\rho_g(X_{t-}) + \rho_g(X_t)}{2}$$

as $n, N \to \infty$. Convergence is uniform in t, if $t \mapsto X_t$ is continuous.

Proof. Similar as in [1] or [5, Section 4.4.1] and the proof of Theorem 5.3, we strengthen assumption (H) together with Condition (5.3) to boundedness of $b^Y(\chi')$ and X.

Let us decompose $\left(\widehat{\rho_g(X)}_t^{n,N} - \frac{\rho_g(X_{t-})+\rho_g(X_t)}{2}\right)$ into

$$\begin{aligned}
&\left\| \frac{1}{T}\sum_{k=-N}^{N}\left(\left(1-\frac{|k|}{N}\right)e^{\mathrm{i}\frac{2\pi}{T}kt}V(Y,g,k)_T^n\right) - \frac{\rho_g(X_{t-})+\rho_g(X_t)}{2}\right\| \\
\le\quad & \frac{1}{T}\left\|\sum_{k=-N}^{N}\left(1-\frac{|k|}{N}\right)e^{\mathrm{i}\frac{2\pi}{T}kt}\left(V(Y,g,k)_T^n - T\mathcal{F}(\rho_g(X))(k)\right)\right\| \\
&+\left\|\sum_{k=-N}^{N}\left(1-\frac{|k|}{N}\right)e^{\mathrm{i}\frac{2\pi}{T}kt}\mathcal{F}(\rho_g(X))(k) - \frac{\rho_g(X_{t-})+\rho_g(X_t)}{2}\right\|.
\end{aligned}$$

By Féjer's theorem the second term converges almost surely to 0. If $t \mapsto X_t$ is continuous, then convergence holds uniformly in t. For the first term, let us decompose $\sum_{k=-N}^{N}\left(1-\frac{|k|}{N}\right)e^{\mathrm{i}\frac{2\pi}{T}kt}(V(Y,g,k)_T^n - T\mathcal{F}(\rho_g(X))(k))$ into

$$\begin{aligned}
&\sum_{k=-N}^{N}\left(1-\frac{|k|}{N}\right)e^{\mathrm{i}\frac{2\pi}{T}kt}\left(V(Y,g,k)_T^n - T\mathcal{F}(\rho_g(X))(k)\right) \\
&=\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)\left(g(\sqrt{n}\Delta_m^n Y) - g(\beta_m^n)\right) \\
&\quad+\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)\left(g(\beta_m^n) - \rho_{m-1}^n(g)\right) \\
&\quad+\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)\rho_{m-1}^n(g) - \int_0^T F_N\left(\frac{2\pi}{T}(t-s)\right)\rho_g(X_s)ds,
\end{aligned}$$

where

$$F_N(x) = \sum_{k=-N}^{N}\left(1-\frac{|k|}{N}\right)e^{\mathrm{i}kx}, \quad \beta_m^n = \sqrt{n}\sqrt{X_{t_{m-1}^n}}\Delta_m^n Z, \quad \rho_m^n(g) = \rho_g(X_{t_m^n}).$$

Since $t \to \rho_g(X_t)$ is càdlàg and bounded (recall the boundedness condition on X and the fact that g has at most polynomial growth) and $\gamma > 1$, the last term in the above sum converges almost surely to 0, uniformly in t, by Riemann integrability.

As a consequence of the proof of Theorem 6.6 below, the second term converges to 0 in probability, uniformly in t, under the assumptions (H) and (J).

Finally we have to focus on the first term, which we decompose into

$$\begin{aligned}
&\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)\left(g(\sqrt{n}\Delta_m^n Y) - g(\sqrt{n}\Delta_m^n D^Y(\chi'))\right) \\
&\quad+\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)\left(g(\sqrt{n}\Delta_m^n D^Y(\chi')) - g(\beta_m^n)\right),
\end{aligned}$$

where $D^Y(\chi')$ is defined in (5.2). The second term converges in probability to 0, uniformly in t, since it can be estimated by

$$\sup_m \mathbb{E}\left[\|g(\sqrt{n}\Delta_m^n D^Y(\chi')) - g(\beta_m^n)\|\right] \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N\left(\frac{2\pi}{T}(t - t_{m-1}^n)\right)$$

and we have $\sup_m \mathbb{E}\left[\|g(\sqrt{n}\Delta_m^n D^Y(\chi')) - g(\beta_m^n)\|\right] \to 0$ (see the proof of [1, Lemma 5.3, Lemma 5.4]). Writing $\zeta_m^n = (g(\sqrt{n}\Delta_m^n Y) - g(\sqrt{n}\Delta_m^n D^Y(\chi')))$ and $\zeta_m^n(p) = (g(\sqrt{n}\Delta_m^n Y(p)) - g(\sqrt{n}\Delta_m^n D^Y(p)(\chi')))$, where $Y(p)$ is defined in (5.4), we have

$$\begin{aligned}(6.7)\qquad & \mathbb{P}\left[\left\|\sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N\left(\frac{2\pi}{T}(t - t_{m-1}^n)\right)\zeta_m^n\right\| > \varepsilon\right] \\ & \leq \mathbb{P}\left[\tau_p \leq T\right] + \mathbb{P}\left[\left\|\sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N\left(\frac{2\pi}{T}(t - t_{m-1}^n)\right)\zeta_m^n(p)\right\| > \varepsilon\right].\end{aligned}$$

Since $\mathbb{P}\left[\tau_p \leq T\right] \to 0$ as $p \to \infty$ and since $\mathbb{E}\left[\| \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N\left(\frac{2\pi}{T}(t - t_{m-1}^n)\right)\zeta_m^n(p)\|\right]$ can be estimated by

$$\sup_m \mathbb{E}\left[\|\zeta_m^n(p)\|\right] \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N\left(\frac{2\pi}{T}(t - t_{m-1}^n)\right),$$

which converges to 0 for all p due to Assumption $(L(0))$, (6.7) tends to 0 as well. ☐

The following corollary states explicit conditions on g and the jumps of Y such that $(L(0))$ is satisfied and relies on Proposition 5.6 (i) above.

Corollary 6.4. *Let g be continuous with $g(x) = o(\|x^2\|)$ as $\|x\| \to \infty$. Let $\gamma > 1$ and suppose that $\lim \frac{n}{N^\gamma} = K$ for some constant $K > 0$. Then under assumption (H) we have for every $t \in [0, T]$*

$$\widehat{\rho_g(X)}_t^{n,N} \xrightarrow{\mathbb{P}} \frac{\rho_g(X_{t-}) + \rho_g(X_t)}{2}$$

as $n, N \to \infty$. Convergence is uniform in t, if $t \mapsto X_t$ is continuous.

Proof. The proof is a consequence of Theorem 6.3 and Proposition 5.6 (i) (as in Corollary 5.8 (i) above). ☐

As mentioned in the introduction of this section, we can now transfer the consistency result to instantaneous covariance estimator (6.2).

Corollary 6.5. *Let g be such that $\rho_g(x) : S_d \times S_d$, $x \mapsto \rho_g(x)$ has a continuous inverse. Then under the assumptions of Theorem 6.3 or Corollary 6.4 we have for every $t \in [0, T]$*

$$\widehat{X}_t^{n,N} \xrightarrow{\mathbb{P}} \rho_g^{-1}\left(\frac{\rho_g(X_{t-}) + \rho_g(X_t)}{2}\right)$$

as $n, N \to \infty$. If $t \mapsto X_t$ is continuous, then

$$\widehat{X}_t^{n,N} \xrightarrow{\mathbb{P}} X_t$$

uniformly in t.

Proof. This corollary is simply a consequence of the continuous mapping theorem. ☐

6.2. **Central limit theorem.** This section is dedicated to the proof of the central limit theorem for (functions of) the instantaneous covariance. Here we need to assume that the trajectories of X are almost surely Hölder continuous with some exponent δ, which thus strengthens assumption $(H1)$ and excludes covariance processes exhibiting jumps.

Theorem 6.6. *Assume that the paths of X are almost surely Hölder continuous with exponent δ. Let $1 < \gamma < 2\delta + 1$ and suppose that $\lim \frac{n}{N^\gamma} = K$ for some constant $K > 0$. Then under $(H1)$ and (K) or (K') and $(L(\eta))$ with $\eta \geq \frac{\gamma-1}{2\gamma}$, the random variable*

$$n^{\frac{\gamma-1}{2\gamma}} \left(\widehat{\rho_g(X)}_t^{n,N} - \rho_g(X_t) \right) \tag{6.8}$$

converges for each $t \in [0,T]$ as $n, N \to \infty$ stably in law to an $\mathcal{F}$-conditional Gaussian random variable defined on an extension of the original probability space with mean 0 and finite non-zero covariance function given by

$$\begin{aligned} V_t^{iji'j'} &:= \lim_{N \to \infty} \frac{1}{T^2 K^{\frac{1}{\gamma}}} \int_0^T (\rho_{g_{ij} g_{i'j'}}(X_s) - \rho_{g_{ij}}(X_s)\rho_{g_{i'j'}}(X_s)) \frac{F_N^2\left(\frac{2\pi}{T}(t-s)\right)}{N} ds \\ &= \frac{2}{3TK^{\frac{1}{\gamma}}} (\rho_{g_{ij} g_{i'j'}}(X_t) - \rho_{g_{ij}}(X_t)\rho_{g_{i'j'}}(X_t)). \end{aligned}$$

Remark 6.7. As already mentioned at the beginning of the article, it is remarkable that the asymptotic variance of the Fourier-Féjer estimator is smaller that the one of the classical local realized variance estimator. For simplicity, let us consider the one-dimensional case with $g(y) = y^2$. Then the variance of the Fourier-Féjer estimator equals

$$\frac{2}{3} \frac{2X_t^2}{TK^{\frac{1}{\gamma}}}. \tag{6.9}$$

under the above assumptions. In comparison, consider the classical (non-truncated) local realized variance estimator given by

$$\widehat{\widehat{X}}_t^{n,N} = \sum_{j=1}^{N} \varepsilon_j^N(t) \sum_{m=1}^{\lfloor nT \rfloor} (\Delta_m^n Y)^2 \varepsilon_j^N(t_{m-1}^n)$$

with

$$\varepsilon_j^N(t) = 1_{[t_{j-1}^N, t_j^N]}(t) \frac{1}{\sqrt{t_j^N - t_{j-1}^N}} \quad \text{and} \quad t_j^N = \frac{jT}{N}, \, j = 1, \ldots, N.$$

Similar as in the above theorem, suppose $1 < \gamma < 2$ and $\lim \frac{n}{N^\gamma} = K$ for some constant $K > 0$. Then, according to [5, Theorem 13.3.3 b)]

$$n^{\frac{\gamma-1}{2\gamma}} \left(\widehat{\widehat{X}}_t^{n,N} - X_t \right)$$

converges for each $t \in [0,T]$ as $n, N \to \infty$ stably in law to an $\mathcal{F}$-conditional Gaussian random variable with mean 0 and covariance function given by

$$\frac{2X_t^2}{TK^{\frac{1}{\gamma}}}.$$

and is therefore $\frac{3}{2}$ times bigger than (6.9). Let us remark that in the notation of [5, Theorem 13.3.3 b)], k_n corresponds to $\frac{nT}{N}$, that is, the number of points in the interval $[t^N_{j-1}, t^N_j]$, $\tau = \frac{\gamma-1}{\gamma} \in (0, \frac{1}{2})$ and $\beta' = K^{\frac{1}{\gamma}} T$ such that the above assertion concerning the classical estimator is implied by [5, Theorem 13.3.3 a)] with $\beta = 0$ since $\tau < \frac{1}{2}$.

Proof. Similarly as in the proof of Theorem 5.3, we strengthen the assumption $(H1)$ to $(SH1)$, that is, $b^Y(\chi')$, defined in (5.3), X and the differential characteristics of the drift and the diffusion part of $\sqrt{X}$ and (3.2) are bounded by a constant. Analogously to the proof of Theorem 6.3 we now decompose

$$n^{\frac{\gamma-1}{2\gamma}} \left(\widehat{\rho_g(X)}^{n,N}_t - \rho_g(X) \right)$$

into

$$\begin{aligned}
& n^{\frac{\gamma-1}{2\gamma}} \left(\frac{1}{T} \sum_{k=-N}^{N} \left(\left(1 - \frac{|k|}{N} \right) e^{\mathrm{i}\frac{2\pi}{T}kt} V(Y,g,k)^n_T \right) - \rho_g(X_t) \right) \\
\leq \quad & n^{\frac{\gamma-1}{2\gamma}} \frac{1}{T} \left(\sum_{k=-N}^{N} \left(1 - \frac{|k|}{N} \right) e^{\mathrm{i}\frac{2\pi}{T}kt} \left(V(Y,g,k)^n_T - T\mathcal{F}(\rho_g(X))(k) \right) \right) + \\
& + n^{\frac{\gamma-1}{2\gamma}} \left| \sum_{k=-N}^{N} \left(1 - \frac{|k|}{N} \right) e^{\mathrm{i}\frac{2\pi}{T}kt} \mathcal{F}(\rho_g(X))(k) - \rho_g(X_t) \right|.
\end{aligned}$$

Due to the Hölder continuity assumption on $t \mapsto X_t$, the second term can be estimated by $Cn^{\frac{\gamma-1}{2\gamma}} N^{-\delta}$ (compare [9, Eq. 13]). Since $\delta > \frac{\gamma-1}{2}$ by assumption, the second term converges almost surely to 0, uniformly in t.

The first term is decomposed as before into

$$\begin{aligned}
& n^{\frac{\gamma-1}{2\gamma}} \frac{1}{T} \sum_{k=-N}^{N} \left(1 - \frac{|k|}{N} \right) e^{\mathrm{i}\frac{2\pi}{T}kt} \left(V(Y,g,k)^n_T - T\mathcal{F}(\rho_g(X))(k) \right) \\
& = n^{\frac{\gamma-1}{2\gamma}} \frac{1}{T} \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N \left(\frac{2\pi}{T}(t - t^n_{m-1}) \right) \left(g(\sqrt{n}\Delta^n_m Y) - g(\beta^n_m) \right) + \\
& + n^{\frac{\gamma-1}{2\gamma}} \frac{1}{T} \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N \left(\frac{2\pi}{T}(t - t^n_{m-1}) \right) \left(g(\beta^n_m) - \rho^n_{m-1}(g) \right) + \\
& + n^{\frac{\gamma-1}{2\gamma}} \frac{1}{T} \left(\sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N \left(\frac{2\pi}{T}(t - t^n_{m-1}) \right) \rho^n_{m-1}(g) - \right. \\
& \qquad \left. - \int_0^T F_N \left(\frac{2\pi}{T}(t-s) \right) \rho_g(X_s) ds \right).
\end{aligned}$$

Due to Lemma 6.1, the last term of this sum can be estimated by

$$Cn^{\frac{\gamma-1}{2\gamma}} \max \left(\frac{1}{N^{\gamma-1}}, \frac{1}{N^{\delta\gamma}} \right) = C \left(\frac{n}{N^\gamma} \right)^{\frac{\gamma-1}{2\gamma}} N^{\frac{\gamma-1}{2}} \max \left(\frac{1}{N^{\gamma-1}}, \frac{1}{N^{\delta\gamma}} \right),$$

which converges to 0, since $\frac{\gamma-1}{2} < \min(\gamma-1, \delta\gamma)$ as a consequence of the assumption $\delta > \frac{\gamma-1}{2}$.

Let us now turn to the second term

$$\begin{aligned}&\frac{1}{T}n^{\frac{\gamma-1}{2\gamma}}\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)(g(\beta^n_m)-\rho^n_{m-1}(g))\\ &\quad=\sum_{m=1}^{\lfloor nT\rfloor}Z^{n,N}_m,\end{aligned}$$

where

$$Z^{n,N}_m=\frac{1}{T}\frac{1}{n^{\frac{\gamma+1}{2\gamma}}}F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)(g(\beta^n_m)-\rho^n_{m-1}(g))$$

Since $\mathbb{E}\left[(g(\beta^n_m)-\rho^n_{m-1}(g))|\mathcal{F}_{t^n_{m-1}}\right]=0$, we also have

$$\lim_{n,N\to\infty}\sum_{m=1}^{\lfloor nT\rfloor}\mathbb{E}\left[Z^{n,N}_m|\mathcal{F}_{t^n_{m-1}}\right]=0.$$

Moreover,

$$\begin{aligned}&\mathbb{E}\left[Z^{n,N}_{m,ij}\overline{Z}^{n,N}_{m,i'j'}|\mathcal{F}_{t^n_{m-1}}\right]\\ &\quad=\frac{1}{T^2}(\rho^n_{m-1}(g_{ij}g_{i'j'})-\rho^n_{m-1}(g_{ij})\rho^n_{m-1}(g_{i'j'}))\frac{1}{n^{\frac{\gamma+1}{\gamma}}}F^2_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right).\end{aligned}$$

Thus we have

$$\begin{aligned}&\sum_{m=1}^{\lfloor nT\rfloor}\mathbb{E}\left[Z^{n,N}_{m,ij}\overline{Z}^{n,N}_{m,i'j'}|\mathcal{F}_{t^n_{m-1}}\right]\\ &\quad=\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{T^2}(\rho^n_{m-1}(g_{ij}g_{i'j'})-\rho^n_{m-1}(g_{ij})\rho^n_{m-1}(g_{i'j'}))\frac{1}{n^{\frac{\gamma+1}{\gamma}}}F^2_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)\\ &\quad=\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{T^2}(\rho^n_{m-1}(g_{ij}g_{i'j'})-\rho^n_{m-1}(g_{ij})\rho^n_{m-1}(g_{i'j'}))\frac{1}{n}\frac{1}{K^{\frac{1}{\gamma}}}\frac{F^2_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)}{N}.\end{aligned}$$

Due to Lemma 6.1 and Remark 6.2 the limit of this expression is given by

$$\begin{aligned}V^{iji'j'}_t&:=\lim_{N\to\infty}\frac{1}{T^2K^{\frac{1}{\gamma}}}\int_0^T(\rho_{g_{ij}g_{i'j'}}(X_s)-\rho_{g_{ij}}(X_s)\rho_{g_{i'j'}}(X_s))\frac{F^2_N\left(\frac{2\pi}{T}(t-s)\right)}{N}ds\\ &=\frac{2}{3TK^{\frac{1}{\gamma}}}(\rho_{g_{ij}g_{i'j'}}(X_t)-\rho_{g_{ij}}(X_t)\rho_{g_{i'j'}}(X_t))\end{aligned}$$

In view of Theorem [6, Theorem IX.7.28] it remains to verify that

$$\sum_{m=1}^{\lfloor nT\rfloor}\mathbb{E}\left[\|Z^{n,N}_m\|^2 1_{\{\|Z^{n,N}_m\|>\varepsilon\}}|\mathcal{F}_{t^n_{m-1}}\right]\xrightarrow{\mathbb{P}}0 \tag{6.10}$$

for all $\varepsilon>0$. By the Cauchy-Schwarz inequality we have

$$\mathbb{E}\left[\|Z^{n,N}_m\|^2 1_{\{|Z^{n,N}_m|>\varepsilon\}}|\mathcal{F}_{t^n_{m-1}}\right]\le\sqrt{\mathbb{E}\left[\|Z^{n,N}_m\|^4|\mathcal{F}_{t^n_{m-1}}\right]}\sqrt{\mathbb{E}\left[1_{\{|Z^{n,N}_m|>\varepsilon\}}|\mathcal{F}_{t^n_{m-1}}\right]}.$$

By definition of $Z_m^{n,N}$ and the polynomial growth of g, we can further estimate

$$\begin{aligned}
&\sqrt{\mathbb{E}\left[\|Z_m^{n,N}\|^4|\mathcal{F}_{t_{m-1}^n}\right]}\\
=&\sqrt{\mathbb{E}\left[\frac{1}{T^4}\left\|g\left(\beta_m^n\right)-\rho_{m-1}^n(g)\right\|^4\frac{1}{n^{\frac{2(\gamma+1)}{\gamma}}}F_N^4\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)|\mathcal{F}_{t_{m-1}^n}\right]}\\
\leq& C\frac{1}{T^2}\frac{1}{n^{\frac{(\gamma+1)}{\gamma}}}F_N^2\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right).
\end{aligned}$$

Taking again the polynomial growth of g into account, there exists some $p \geq 0$ such that

$$\|g\left(\beta_m^n\right)-\rho_{m-1}^n(g)\| \leq C(1+\|U_{m-1}^n\|^p), \quad \mathbb{P}\text{-a.s.},$$

where $U_{m-1}^n=\sqrt{n}\sqrt{X_{t_{m-1}^n}}\Delta_m^n Z \sim N(0,X_{t_{m-1}^n})$. Thus

$$\begin{aligned}
&\mathbb{E}\left[1_{\{\|Z_m^{n,N}\|>\varepsilon\}}|\mathcal{F}_{t_{m-1}^n}\right]\\
=&\mathbb{P}\left[\|Z_m^{n,N}\|>\varepsilon|\mathcal{F}_{t_{m-1}^n}\right]\\
\leq&\mathbb{P}\left[C(1+\|U_{m-1}^n\|^p)\frac{1}{n^{\frac{(\gamma+1)}{2\gamma}}}F_N\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)>\varepsilon|\mathcal{F}_{t_{m-1}^n}\right]\\
=&\mathbb{P}\left[\|U_{m-1}^n\|>\underbrace{\left(\frac{1}{C}\left(n^{\frac{(\gamma+1)}{2\gamma}}F_N^{-1}\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)\varepsilon-1\right)\right)^{\frac{1}{p}}}_{\to\infty \text{ as } n,N\to\infty}|\mathcal{F}_{t_{m-1}^n}\right].
\end{aligned}$$

Since this tends to 0, we can estimate (6.10) by

$$\begin{aligned}
&\sum_{m=1}^{\lfloor nT\rfloor}C\frac{1}{T^2}\frac{1}{n^{\frac{(\gamma+1)}{\gamma}}}F_N^2\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)\sqrt{\mathbb{P}\left[\|Z_m^{n,N}\|>\varepsilon|\mathcal{F}_{t_{m-1}^n}\right]}\\
=&\sum_{m=1}^{\lfloor nT\rfloor}C\frac{1}{T^2}\frac{1}{n}\left(\frac{N^\gamma}{n}\right)^{\frac{1}{\gamma}}\frac{F_N^2\left(\frac{2\pi}{T}(t-t_{m-1}^n)\right)}{N}\sqrt{\mathbb{P}\left[\|Z_m^{n,N}\|>\varepsilon|\mathcal{F}_{t_{m-1}^n}\right]}\xrightarrow{\mathbb{P}}0,
\end{aligned}$$

where convergence to 0 follows from Lemma 6.1 and the above estimate for

$$\mathbb{P}\left[\|Z_m^{n,N}\|>\varepsilon|\mathcal{F}_{t_{m-1}^n}\right],$$

hence Equation (6.10) is verified. Moreover, similarly as in the proof of Proposition 5.5, we have

$$\mathbb{E}\left[Z_m^{n,N}\Delta_m^n Z|\mathcal{F}_{t_{m-1}^n}\right]=0$$

and

$$\mathbb{E}\left[Z_m^{n,N}\Delta_m^n M|\mathcal{F}_{t_{m-1}^n}\right]=0$$

for any bounded martingale M which is orthogonal to the Brownian motion Z.

Finally we have to consider

$$
\begin{aligned}
n^{\frac{\gamma-1}{2\gamma}} \frac{1}{T} \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)(g(\sqrt{n}\Delta^n_m Y)-g(\beta^n_m))\\
= n^{\frac{\gamma-1}{2\gamma}} \frac{1}{T} \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)(g(\sqrt{n}\Delta^n_m Y)-g(\sqrt{n}\Delta^n_m D^Y(\chi')))+\\
+ n^{\frac{\gamma-1}{2\gamma}} \frac{1}{T} \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)(g(\sqrt{n}\Delta^n_m D^Y(\chi'))-g(\beta^n_m)).
\end{aligned}
$$

In view of Lemma [5, Lemma 2.2.11] it is sufficient to prove that

$$
\frac{1}{T^2} \sum_{m=1}^{\lfloor nT \rfloor} \mathbb{E}\left[\|U^{n,1}_m\|^2\right] \to 0, \tag{6.11}
$$

$$
\frac{1}{T} \sum_{m=1}^{\lfloor nT \rfloor} \mathbb{E}\left[U^{n,1}_m | \mathcal{F}_{t^n_{m-1}}\right] \xrightarrow{\mathbb{P}} 0, \tag{6.12}
$$

and

$$
\frac{1}{T}\mathbb{P}\left[\left\|\sum_{m=1}^{\lfloor nT \rfloor} U^{n,2}_m\right\| > \varepsilon\right] \to 0\,, \tag{6.13}
$$

where

$$
\begin{aligned}
U^{n,1}_m &= \frac{1}{n^{\frac{\gamma+1}{2\gamma}}} F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)(g(\sqrt{n}\Delta^n_m D^Y(\chi'))-g(\beta^n_m))\,,\\
U^{n,2}_m &= \frac{1}{n^{\frac{\gamma+1}{2\gamma}}} F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)(g(\sqrt{n}\Delta^n_m Y)-g(\sqrt{n}\Delta^n_m D^Y(\chi')).
\end{aligned}
$$

Let us first focus on $U^{n,1}_m$. By [1, Lemma 5.3 and Lemma 5.4]) and the Hölder continuity of X we have

$$
\sup_m \mathbb{E}\left[\left\|g(\sqrt{n}\Delta^n_m D^Y(\chi'))-g(\beta^n_m))\right\|^2\right] \to 0
$$

and we can therefore estimate (6.11) by

$$
\sup_m \mathbb{E}\left[\left\|g(\sqrt{n}\Delta^n_m D(\chi')-g(\beta^n_m))\right\|^2\right] \frac{1}{T} \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n^{\frac{\gamma+1}{\gamma}}} F^2_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right),
$$

which converges to 0 due to Lemma 6.1.

Concerning (6.12), it is possible to decompose

$$
g(\sqrt{n}\Delta^n_m D^Y(\chi'))-g(\beta^n_m) = A^n_m + B^n_m,
$$

where for all m, $\mathbb{E}\left[A^n_m|\mathcal{F}_{t^n_{m-1}}\right]=0$ and $\mathbb{E}\left[\|B^n_m\|\right]\leq\frac{1}{n^{\eta}}$ with $\eta>\frac{\gamma-1}{2\gamma}$ (see [5, Section 5.3.3, C]. Then

$$
\begin{aligned}
&n^{\frac{\gamma-1}{2\gamma}}\frac{1}{T}\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)\mathbb{E}\left[g(\sqrt{n}\Delta^n_m D^Y(\chi'))-g(\beta^n_m)|\mathcal{F}_{t^n_{m-1}}\right]\\
&=n^{\frac{\gamma-1}{2\gamma}}\frac{1}{T}\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right)\mathbb{E}\left[A^n_m+B^n_m|\mathcal{F}_{t^n_{m-1}}\right]\\
&\leq n^{\frac{\gamma-1}{2\gamma}}\sup_m\mathbb{E}\left[\|B^n_m\|\right]\frac{1}{T}\sum_{m=1}^{\lfloor nT\rfloor}\frac{1}{n}F_N\left(\frac{2\pi}{T}(t-t^n_{m-1})\right),
\end{aligned}
$$

converges 0 due to Lemma 6.1 and thus yields (6.12). Condition (6.13) follows from the assumption $L(\eta)$ for $\eta\geq\frac{\gamma-1}{2\gamma}$ and a similar estimate as in (6.7). The assertion is now implied by all these estimates and [6, Theorem IX.7.28]. $\square$

Remark 6.8.

(i) In the above theorem the assumption that the trajectories of X are δ-Hölder continuous is crucial. Typically this can be verified by using Kolmogorov's criterium, which states that the Hölder exponent δ satisfies $\delta<\frac{\varepsilon}{\alpha}$ if

$$\mathbb{E}\left[\|X_t-X_s\|^{\alpha}\right]\leq C|t-s|^{1+\varepsilon}.$$

In our case we have for $\alpha>2$

$$
\begin{aligned}
&\mathbb{E}\left[\left\|\int_s^t b^X_u du+\sum_{j=1}^p\int_s^t Q^j_u dB_{u,j}\right\|^{\alpha}\right]\\
\leq &C\left(\mathbb{E}\left[\left\|\int_s^t b^X_u du\right\|^{\alpha}\right]+\sum_{j=1}^p\mathbb{E}\left[\left\|\int_s^t Q^j_u dB_{u,j}\right\|^{\alpha}\right]\right)\\
\leq &C\left(\mathbb{E}\left[\int_s^t |t-s|^{\alpha-1}\|b^X_u\|^{\alpha}du\right]+\sum_{j=1}^p\mathbb{E}\left[\left\|\int_s^t (Q^j_u)^2 du\right\|^{\frac{\alpha}{2}}\right]\right)\\
\leq &C|t-s|^{\alpha}+C\sum_{j=1}^p\mathbb{E}\left[\int_s^t |t-s|^{\frac{\alpha}{2}-1}\|Q^j_u\|^{\alpha}du\right]\\
\leq &C|t-s|^{\alpha}+C|t-s|^{\frac{\alpha}{2}}\\
\leq &C|t-s|^{\frac{\alpha}{2}}
\end{aligned}
$$

provided that $\mathbb{E}\left[\|b^X_u\|^{\alpha}\right]<\infty$ and $\mathbb{E}\left[\|Q^j_u\|^{\alpha}\right]<\infty$. For the last inequality we used the fact that we are working on $[0,T]$. If these moments exist for all α then we have Hölder continuity with $\delta<\frac{1}{2}$. This is for example satisfied for affine diffusion processes on S^+_d.

(ii) If we have Hölder continuity for all $\delta<\frac{1}{2}$, then $\gamma\in(1,2)$. The higher γ the better the convergence rate and it lies between $(0,\frac{1}{4})$.

(iii) Condition $(L(\eta))$ with $\eta\geq\frac{\gamma-1}{2\gamma}$ is satisfied, if the assumptions of Proposition 5.6 (ii) hold with $r>\frac{\gamma-1}{\gamma}\frac{\beta}{2-\beta}$ and $\beta<1+\frac{1}{2\gamma-1}$.

Similarly as above for the consistency statement we now translate the central limit theorem to $\widehat{X}_t^{n,N}$ defined in (6.2).

Corollary 6.9. *Let g be such that $\rho_g(x) : S_d \times S_d$, $x \mapsto \rho_g(x)$ has a differentiable inverse. Then under the assumptions of Theorem 6.6*

$$n^{\frac{\gamma-1}{2\gamma}} \left(\widehat{X}_t^{n,N} - X_t \right)$$

converges as $n, N \to \infty$ for each $t \in [0, T]$ stably in law to a $\mathcal{F}$-conditional Gaussian random variable given by

$$M_t = (\nabla \rho_g(X_t))^{-1} N_t,$$

where N_t denotes the limit of (6.8). In particular the covariance of M_t is given by

$$U_t^{iji'j'} = \sum_{k,l,k',l'} (\nabla \rho_g(X_t))^{-1}_{ijkl} (\nabla \rho_g(X_t))^{-1}_{i'j'k'l'} V_t^{klk'l'}.$$

Proof. This assertion follows from the concept of stable convergence and is known as generalized Δ-method (see [11, Theorem 1.10]). □

7. Covariance of covariance estimation

Having reconstructed the path of the instantaneous covariance, we can now proceed with step (3), that is, the estimation of functions of the integrated covariance. To this end we plug the reconstructed path of the instantaneous covariance process into jump robust estimators as introduced in Section 5:

$$V(\widehat{X}^{n,N}, f, 0)_T^m := \frac{1}{m} \sum_{p=1}^{\lfloor mT \rfloor} f(\sqrt{m} \Delta_p^m \widehat{X}^{n,N}),$$

where $f : S_d \to \mathbb{R}^{d^2 \times d^2}$ satisfies certain properties which are specified in Theorem 7.1 below. In order to formulate the following theorem, we denote by X^D the diffusion part of X, that is,

$$X_t^D = x + \int_0^t b_s^X ds + \sum_{j=1}^p \int_0^t Q_{s-}^j dB_{s,j}$$

and by Y^D we mean

$$Y_t^D = y + \int_0^t b_s^Y ds + \int_0^t \sqrt{X_s^D} dZ_s + \int_0^t \int_{\mathbb{R}^d} \chi(\xi)(\mu^Y(d\xi, ds) - K_s(d\xi)ds) + \int_0^t \int_{\mathbb{R}^d} \xi \mu^Y(d\xi, ds).$$

Moreover,

$$\widehat{\rho_g(X^D)}_t^{n,N} := \frac{1}{T} \sum_{k=-N}^{N} \left(1 - \frac{|k|}{N}\right) e^{\mathrm{i}\frac{2\pi}{T}kt} V(Y^D, g, k)_T^n$$

and

$$\widehat{(X^D)}_t^{n,N} := \rho_g^{-1} \left(\widehat{\rho_g(X^D)}_t^{n,N} \right).$$

Theorem 7.1.

- *Assume that X has jumps are of finite activity and suppose that the paths of X^D are almost surely Hölder continuous with exponent δ.*

- *Let* $1 < \gamma < 2\delta + 1$ *and suppose that* $\lim \frac{n}{N^\gamma} = K$ *for some constant* $K > 0$.
- *Let the conditions* $(H1)$ *and* (K) *or* (K') *and* $L(\eta)$ *with* $\eta \geq \frac{\gamma-1}{2\gamma}$ *be in force and suppose that the instantaneous covariance process* q *of* X *defined by* $q_{s,iji'j'} = \sum_{k,l} Q^k_{s,ij} Q^l_{s,i'j'}$ *satisfies* $(H1)$.
- *Let* g *be such that* $x \mapsto (\nabla \rho_g(x))^{-1}$ *exists and is continuous.*
- *Suppose that* $f : S_d \to \mathbb{R}^{d^2 \times d^2}$ *satisfies either* (K) *or* (K') *and is additionally globally* α*-Hölder continuous for* $\alpha \in (0,1)$.
- *Let* $\iota < \frac{\gamma-1}{\gamma}\frac{\alpha}{1+\alpha}$ *and assume that* $\lim \frac{m}{n^\iota} = \widetilde{K}$ *for some constant* $\widetilde{K} > 0$.

Then

$$n^{\frac{\iota}{2}} \left(V(\widehat{X}^{n,N}, f, 0)^m_T - \int_0^T \rho_f(q_s) ds \right)$$

converges as $n, N, m \to \infty$ *stably in law to a* $\mathcal{F}$*-conditional Gaussian random variable defined on an extension of the original probability space with* 0 *mean and covariance function given by*

$$F_{ijkli'j'k'l'} = \frac{1}{\widetilde{K}} \int_0^T \left(\rho_{f_{iji'j'} f_{klk'l'}}(q_s) + \rho_{f_{ijk'l'}}(q_s) \rho_{f_{i'j'kl}}(q_s) \right) ds.$$

Proof. Let us decompose

$$\sqrt{m} \left(V(\widehat{X}^{n,N}, f, 0)^m_T - \int_0^T \rho_f(q_s) ds \right) = \sqrt{m} \left(V(\widehat{X}^{n,N}, f, 0)^m_T - V(X^D, f, 0)^m_T \right) + \tag{7.1}$$
$$+ \sqrt{m} \left(V(X^D, f, 0)^m_T - \int_0^T \rho_f(q_s) ds) \right).$$

In view [5, Theorem 5.3.5 and 5.3.6] and the relation between m and n, the second term converges to the stated Gaussian random variable. Hence we only have to prove that the first term converges to 0 in probability.

Decomposing $\widehat{X}^{n,N}_t$ into

$$\widehat{X}^{n,N}_t = \widehat{(X^D)}^{n,N}_t + \left(\widehat{X}^{n,N}_t - \widehat{(X^D)}^{n,N}_t \right),$$

the first term of the right hand side of (7.1) can be estimated due to the assumptions on f by

$$\begin{aligned}
&\sqrt{m} \left\| V(\widehat{X}^{n,N}, f, 0)^m_T - V(X^D, f, 0)^m_T \right\| \\
&\leq \frac{1}{\sqrt{m}} \sum_{p=1}^{\lfloor mT \rfloor} \left\| f\left(\sqrt{m} \Delta^m_p \widehat{X}^{n,N} \right) - f\left(\sqrt{m} \Delta^m_p X^D \right) \right\| \\
&\leq m^{\frac{\alpha-1}{2}} C \sum_{p=1}^{\lfloor mT \rfloor} \left(\left\| \Delta^m_p \widehat{(X^D)}^{n,N} - \Delta^m_p X^D \right\|^\alpha + \left\| \Delta^m_p \widehat{X}^{n,N}_t - \Delta^m_p \widehat{(X^D)}^{n,N}_t \right\|^\alpha \right) \\
&\leq m^{\frac{\alpha+1}{2}} 2C \sup_p \left\| \widehat{(X^D)}^{n,N}_{t^m_p} - X^D_{t^m_p} \right\|^\alpha + m^{\frac{\alpha+1}{2}} 2C \sup_p \left\| \widehat{X}^{n,N}_{t^m_p} - \widehat{(X^D)}^{n,N}_{t^m_p} \right\|^\alpha .
\end{aligned}$$

By Lemma 7.2 and 7.3 below, the relation between m and n and the condition on ι, both summands converge to 0 in probability, which proves the theorem. □

Lemma 7.2. *Let the conditions of Theorem 6.6 be in force and let g be such that $x \mapsto (\nabla \rho_g(x))^{-1}$ exists and is continuous. Moreover, consider a sequence $(t_p)_p$ in $[0,T]$ and let $\alpha \in (0,1]$. Then for every $\theta < \frac{\gamma-1}{2\gamma}$*

$$n^{\alpha\theta} \sup_p \|\widehat{X}^{n,N}_{t_p} - X_{t_p}\|^\alpha \xrightarrow{\mathbb{P}} 0.$$

Proof. By localizing we can assume that X is uniformly bounded. In fact, consider a localizing sequence

$$\tau_k = \inf\{t \geq 0 \,|\, \|X_t\| > k\}, \quad k \in \mathbb{N},$$

and the processes

$$\begin{aligned} Y(k)_t &= y + \int_0^t b^Y_s ds + \int_0^{t\wedge\tau_k} \sqrt{X_s} dZ_s + \int_0^t \int_{\mathbb{R}^d} \chi(\xi)(\mu^Y(d\xi, ds) - K_s(d\xi)ds) \\ &\qquad + \int_0^t \int_{\mathbb{R}^d} (\xi - \chi(\xi))\mu^Y(d\xi, ds), \\ X(k)_t &= X_t 1_{\{t<\tau_k\}}, \end{aligned}$$

where $(X_t(k))_{t\geq 0}$ is uniformly bounded by definition. Moreover, define

$$\widehat{\rho_g(X(k))}^{n,N}_t := \frac{1}{T} \sum_{k=-N}^{N} \left(1 - \frac{|k|}{N}\right) e^{\mathrm{i}\frac{2\pi}{T}kt} V(Y(k), g, k)^n_T$$

and

$$\widehat{X(k)}^{n,N}_t := \rho_g^{-1}\left(\widehat{\rho_g(X(k))}^{n,N}_t\right).$$

Then the left hand side of

$$\begin{aligned} \mathbb{P}\left[n^{\alpha\theta} \sup_p \|\widehat{X}^{n,N}_{t_p} - X_{t_p}\|^\alpha > \varepsilon\right] &\leq \mathbb{P}\left[\tau_k \leq T\right] \\ &\qquad + \mathbb{P}\left[n^{\alpha\theta} \sup_p \|\widehat{X(k)}^{n,N}_{t_p} - X(k)_{t_p}\|^\alpha > \varepsilon\right] \end{aligned}$$

tends to 0, if the second term on the right hand side does. Therefore, we can assume uniform boundedness of X.

Furthermore, by the continuous mapping theorem it suffices to prove

$$n^{\theta} \sup_p \|\widehat{X}^{n,N}_{t_p} - X_{t_p}\| \xrightarrow{\mathbb{P}} 0.$$

By the mean value theorem we obtain the identity

$$n^\theta \sup_p \|\widehat{X}^{n,N}_{t_p} - X_{t_p}\| = n^\theta \sup_p \left\| (\nabla \rho_g(\zeta^{n,N}_{t_p}))^{-1} \left(\widehat{\rho_g(X)}^{n,N}_{t_p} - \rho_g(X_{t_p})\right)\right\|, \tag{7.2}$$

where $\zeta^{n,N}_{t_p}$ is a random variable satisfying $\|\zeta^{n,N}_{t_p} - X_{t_p}\| \leq \|\widehat{X}^{n,N}_{t_p} - X_{t_p}\|$. Due to the continuity assumption on $x \mapsto (\nabla\rho_g(x))^{-1}$ and boundedness of X, (7.2) converges to 0 in probability if

$$n^\theta \sup_p \|\widehat{\rho_g(X)}^{n,N}_{t_p} - \rho_g(X_{t_p})\| \xrightarrow{\mathbb{P}} 0.$$

An inspection of the proof of Theorem 6.6 reveals that this is the case for all $\theta < \frac{\gamma-1}{2\gamma}$. □

Lemma 7.3. *Assume that X has jumps of of finite activity and suppose that the paths of X^D are almost surely Hölder continuous with exponent δ. Let $1 < \gamma < 2\delta+1$ and suppose that $\lim \frac{n}{N^\gamma} = K$ for some constant $K > 0$. Assume that the conditions $(H1)$ and (K) or (K') and $L(\eta)$ with $\eta \geq \frac{\gamma-1}{2\gamma}$ are in force and g be such that $x \mapsto (\nabla \rho_g(x))^{-1}$ exists and is continuous. Consider a sequence $(t_p)_p$ in $[0,T]$ and let $\alpha \in (0,1]$. Then for every $\theta < \frac{\gamma-1}{\gamma}$*

$$n^{\theta\alpha} \sup_p \left\| \widehat{X}^{n,N}_{t_p} - \widehat{(X^D)}^{n,N}_{t_p} \right\|^\alpha \xrightarrow{\mathbb{P}} 0.$$

Proof. By arguing similarly as in Lemma 7.2 above, we can assume that X, X^D and $b^Y(\chi')$ are uniformly bounded by a constant, where $b^Y(\chi')$ is defined in (5.3). The mean value theorem yields

$$\begin{aligned} &n^{\theta\alpha} \sup_p \| \widehat{X}^{n,N}_{t_p} - \widehat{(X^D)}^{n,N}_{t_p} \|^\alpha \\ &\quad = n^{\theta\alpha} \sup_p \left\| (\nabla \rho_g(\zeta^{n,N}_{t_p}))^{-1} \left(\widehat{\rho_g(X)}^{n,N}_{t_p} - \widehat{\rho_g(X^D)}^{n,N}_{t_p} \right) \right\|^\alpha , \end{aligned}$$

where $\zeta^{n,N}_{t_p}$ is a random variable satisfying $\|\zeta^{n,N}_{t_p} - \widehat{(X^D)}^{n,N}_{t_p}\| \leq \|\widehat{X}^{n,N}_{t_p} - \widehat{(X^D)}^{n,N}_{t_p}\|$. Due to the continuity assumption on $x \mapsto (\nabla \rho_g(x))^{-1}$ and boundedness of X, (7.2) converges to 0 in probability if

$$n^{\theta\alpha} \sup_p \left\| \widehat{\rho_g(X)}^{n,N}_{t_p} - \widehat{\rho_g(X^D)}^{n,N}_{t_p} \right\|^\alpha \xrightarrow{\mathbb{P}} 0.$$

By definition of $\widehat{\rho_g(X)}^{n,N}$ and abbreviating $F_N\left(\frac{2\pi}{T}(t_p - t^n_{m-1})\right)$ by F^p_N,

$$\|\widehat{\rho_g(X)}^{n,N}_{t_p} - \widehat{\rho_g(X^D)}^{n,N}_{t_p}\|^\alpha$$

can be written as

$$\begin{aligned} &\sup_p \left\| \sum_{m=1}^{\lfloor nT \rfloor} \frac{1}{n} F^p_N \left(g(\sqrt{n}\Delta^n_m Y) - g(\sqrt{n}\Delta^n_m Y^D) \right) 1_{\{\Delta^n_m Y \neq \Delta^n_m Y^D\}} \right\|^\alpha \\ &\leq \sup_p \sum_{m=1}^{\lfloor nT \rfloor} \left(\frac{1}{n} F^p_N \right)^\alpha \left\| g(\sqrt{n}\Delta^n_m D^Y(\chi')) - g(\sqrt{n}\Delta^n_m D^{Y^D}(\chi')) \right\|^\alpha 1_{\{\Delta^n_m Y \neq \Delta^n_m Y^D\}} \\ &\quad + \sup_p \sum_{m=1}^{\lfloor nT \rfloor} \left(\frac{1}{n} F^p_N \right)^\alpha \left\| g(\sqrt{n}\Delta^n_m Y) - g(\sqrt{n}\Delta^n_m D^Y(\chi')) \right\|^\alpha 1_{\{\Delta^n_m Y \neq \Delta^n_m Y^D\}} \\ &\quad + \sup_p \sum_{m=1}^{\lfloor nT \rfloor} \left(\frac{1}{n} F^p_N \right)^\alpha \left\| g(\sqrt{n}\Delta^n_m D^{Y^D}(\chi')) - g(\sqrt{n}\Delta^n_m Y^D) \right\|^\alpha 1_{\{\Delta^n_m Y \neq \Delta^n_m Y^D\}}. \end{aligned} \tag{7.3}$$

By the boundedness of $b^Y(\chi')$, X and X^D and the polynomial growth of g,

$$\mathbb{E}\left[\left\| g(\sqrt{n}\Delta^n_m D^Y(\chi')) - g(\sqrt{n}\Delta^n_m D^{Y^D}(\chi')) \right\|^\alpha \right]$$

is bounded by a constant. Therefore and due to the finite activity of the jumps of X (note that $\Delta_m^n Y \neq \Delta_m^n Y^D$ only if X jumps)

$$\mathbb{E}\left[\sum_{m=1}^{\lfloor nT \rfloor} \left\| g(\sqrt{n}\Delta_m^n D^Y(\chi')) - g(\sqrt{n}\Delta_m^n D^{Y^D}(\chi')) \right\|^{\alpha} 1_{\{\Delta_m^n Y \neq \Delta_m^n Y^D\}}\right]$$

is bounded as well. This and the form of the Féjer kernel implies that the first summand in (7.3) multiplied with $n^{\theta\alpha}$ can be estimated by

$$Cn^{\theta\alpha}\left(\frac{N}{n}\right)^{\alpha} = \widetilde{C}n^{\theta\alpha}n^{\alpha\frac{1-\gamma}{\gamma}},$$

which converges to 0 due to the condition on θ. As a consequence of $(L(\eta))$, the remaining two summands in (7.3) multiplied with $n^{\theta\alpha}$ converge even faster to 0 and hence the assertion follows. $\square$

8. Simulation results

In this section we illustrate our theoretical results in the case of a multivariate affine model, where both the log-price Y and the instantaneous covariance process X can jump. More precisely, we consider a multivariate Bates-type model of the form

$$\begin{aligned} Y_t &= y + \int_0^t b_s ds + \int_0^t \sqrt{X_{s-}} dZ_s + \int_0^t \int_{\mathbb{R}^d} \xi \mu^Y(d\xi, ds), \\ X_t &= x + \int_0^t (b + MX_t + X_t M^\top)dt + \sqrt{X_t}dB_t\Sigma + \Sigma dB_t^\top \sqrt{X_t} + \\ &\quad + \int_0^t \int_{S_d^+} \xi \mu^X(d\xi, ds), \end{aligned}$$

where

- Z is a d-dimensional Brownian motion correlated with the $d \times d$ matrix of Brownian motions B such that $Z = \sqrt{1 - \rho^\top \rho} W + B\rho$, where $\rho \in [-1, 1]^d$ such that $\rho^\top \rho \leq 1$ and W is a d-dimensional Brownian motion independent of B,
- $\mu^Y(d\xi, dt)$ is the random measure associated with the jumps of Y, whose compensator is given by $\sum_{i=1}^d \lambda^{Y_i} F^{Y_i}(d\xi_i)dt$, where $\lambda^{Y_i} > 0$ and F^{Y_i} denotes the Gaussian density with mean μ_i and standard deviation σ_i,
- $\mu^X(d\xi, dt)$ is the random measure associated with the jumps of X, whose compensator is given by $\lambda^{X_{11}} F^{X_{11}}(d\xi_{11})dt$, where $\lambda^{X_{11}} > 0$ and $F^{X_{11}}$ denotes the density of the exponential distribution with parameter θ,[2]
- the drift of Y is given by $b_{s,i} = -\frac{1}{2} X_{s,ii} - \lambda^{Y_i}(e^{\mu_i - \frac{1}{2}\sigma_i^2} - 1)$ and
- the parameters of X satisfy $M \in \mathbb{R}^{d \times d}$, $\Sigma \in S_d^+$, $b \in S_d^+$ such that

$$b - (d-1)\Sigma^2 \in S_d^+.$$

Note that the truncation function of Y is here chosen to be 0.

As described in Section 1 and Section 4 we aim to recover the instantaneous covariance process X and the parameters $\alpha := \Sigma^2$ and ρ from observations of Y. In order to be in accordance with market specifications, we simulate Y and X on

---

[2] We here only suppose that X_{11} can jump.

$n = 127750 = 511 * 250$ grid points, which corresponds to 1 year ($T = 1$) of 1-minute data. For our numerical simulation, we consider the case $d = 2$ and use the following parameter values:

$(Y_{0,1}, Y_{0,2})$	$(0, 0)$
$\begin{pmatrix} X_{0,11} & X_{0,12} \\ X_{0,12} & X_{0,22} \end{pmatrix}$	$\begin{pmatrix} 0.09 & -0.036 \\ -0.036 & 0.09 \end{pmatrix}$
M	$\begin{pmatrix} -1.6 & -0.2 \\ -0.4 & -1 \end{pmatrix}$
$\alpha = \Sigma^2$	$\begin{pmatrix} 0.0725 & 0.06 \\ 0.06 & 0.1325 \end{pmatrix}$
b	3.5α
ρ	$(-0.3, -0.5)$
$\left(\lambda^{Y_1}, \lambda^{Y_2}\right)$	$(100, 100)$
(μ_1, μ_2)	$(-0.005, -0.003)$
(σ_1, σ_2)	$(0.015, 0.02)$
$\lambda^{X_{11}}$	10
θ	0.05

In order to illustrate in particular that our estimator is robust to small and frequent jumps, the jump intensity of both log-prices is chosen to be quite high. Figure 2 and 3 below show simulated trajectories of the log-price and the instantaneous covariance process, where the jumps are removed in the second graph in each case.

A comparison between the reconstructed and simulated trajectories of the instantaneous covariance process is shown in Figure 4. These figures illustrate that – even in the case of (frequent) jumps in the log-price and in the variance (as it is the case for X_{11}) – the paths of X can be recovered very well. For the reconstruction of the trajectories of X we choose $N = 210$ Fourier coefficients, which corresponds to the choice $\gamma \approx 2$ and $K \approx 3$, as specified in Theorem 6.6. This is a reasonable choice in view of an acceptable bias (in particular due to the fact that we do not sum all Fourier coefficients) and a rather small variance. Both, the simulated as well as the reconstructed trajectories are evaluated at $2N + 1$ points. In our concrete implementation the estimator for the Fourier coefficients (5.1) is based on the Tauchen-Todorov specification of the function g, that is,

$$g : \mathbb{R}^2 \to S_2, \quad (y_1, y_2) \mapsto (\cos(y_i + 1_{\{j \neq i\}} y_j))_{i,j \in \{1,2\}}.$$

In this case

$$\rho_g(x) = \begin{pmatrix} e^{-\frac{1}{2}x_{11}} & e^{-\frac{1}{2}(x_{11}+2x_{12}+x_{22})} \\ e^{-\frac{1}{2}(x_{11}+2x_{12}+x_{22})} & e^{-\frac{1}{2}x_{22}} \end{pmatrix}$$

and $\widehat{X}_t^{n,N}$ is obtained by

$$\widehat{X}_{t,ii}^{n,N} = -2 \log \left(\widehat{\rho_{g_{ii}}(X)}_t^{n,N} \right), \quad i \in \{1, 2\},$$

$$\widehat{X}_{t,12}^{n,N} = \frac{1}{2} \left(-2 \log \left(\widehat{\rho_{g_{ij}}(X)}_t^{n,N} \right) - \widehat{X}_{t,11}^{n,N} - \widehat{X}_{t,22}^{n,N} \right).$$



FIGURE 2. Simulated log-price with and without jumps on $n = 127750$ points

The reconstructed trajectories of X are then used to estimate the parameters α and ρ. To this end, we use the power variation estimators, i.e.,

$$V(\widehat{X}_{ij}^{n,N}, f_r, 0)_1^m := \frac{1}{m} \sum_{p=1}^{m} |\Delta_m^p \widehat{X}_{ij}^{n,N}|^r, \quad i,j \in \{1,2\},$$

$$V(\widehat{X}_{ii}^{n,N}, Y_i, f_{r,s}, 0)_1^m := \frac{1}{m} \sum_{p=1}^{m} |\Delta_m^p \widehat{X}_{ii}^{n,N}|^r |\Delta_m^p Y_i|^s, \quad i \in \{1,2\},$$

where

$$f_r : \mathbb{R} \to \mathbb{R}_+, \quad x \mapsto |x|^r,$$
$$f_{r,s} : \mathbb{R}^2 \to \mathbb{R}_+, \quad (x,y) \mapsto |x|^r |y|^s.$$



FIGURE 3. Simulated instantaneous covariance with and without jumps on $n = 127750$ points

These quantities are estimators for the power (co)variation of X and Y. Indeed we have under the assumptions of Theorem 7.1

$$
\begin{aligned}
V(\widehat{X}_{ii}^{n,N}, f_r, 0)_1^m &\to \int_0^1 \rho_{f_r}(4\alpha_{ii} X_{s,ii}) ds \\
&= \sqrt{\frac{1}{\pi}} 2^{\frac{r}{2}} \Gamma\left(\frac{r+1}{2}\right) (4\alpha_{ii})^{\frac{r}{2}} \int_0^1 X_{s,ii}^{\frac{r}{2}} ds, \\
V(\widehat{X}_{12}^{n,N}, f_r, 0)_1^m &\to \int_0^1 \rho_{f_r}(\alpha_{11} X_{s,11} + 2\alpha_{12} X_{s,12} + \alpha_{22} X_{s,22}) ds \\
&= \sqrt{\frac{1}{\pi}} 2^{\frac{r}{2}} \Gamma\left(\frac{r+1}{2}\right) \int_0^1 (\alpha_{11} X_{s,11} + 2\alpha_{12} X_{s,12} + \alpha_{22} X_{s,22})^{\frac{r}{2}} ds \\
&=: PV_{12}(X, \alpha_{11}, \alpha_{12}, \alpha_{22}), \\
V(\widehat{X}_{ii}^{n,N}, Y_i, f_{r,s}, 0)_1^m &\to \int_0^1 \rho_{f_{r,s}} \left(\begin{pmatrix} 4\alpha_{ii} X_{s,ii} & 2(\sqrt{\alpha}\rho)_i \\ 2(\sqrt{\alpha}\rho)_i X_{s,11} & X_{s,ii} \end{pmatrix} \right) ds \\
&= \frac{1}{\pi} 2^{\frac{r+s}{2}} \Gamma\left(\frac{r+1}{2}\right) \Gamma\left(\frac{s+1}{2}\right) \\
&\quad \times {}_2F_1\left(-\frac{r}{2}, -\frac{s}{2}; \frac{1}{2}; \left(\frac{(\sqrt{\alpha}\rho)_i}{\sqrt{\alpha_{ii}}}\right)^2\right) (4\alpha_{ii})^{\frac{r}{2}} \int_0^1 X_{s,ii}^{\frac{r+s}{2}} ds \\
&=: PC_{ii}(X, Y, \alpha, \rho)
\end{aligned}
$$



Figure 4. Reconstructed and simulated instantaneous covariance evaluated on 421 points

as $m, n, N \to \infty$. The formulas on the right hand sides follow from the expressions for the absolute moments of the bivariate Gaussian distribution (see, e.g.,[10]) and ${}_2F_1\,(a, b; c; x)$ denotes the Gaussian hypergeometric function.

The estimators for α and ρ can now be constructed via

$$
\begin{aligned}
\widehat{\alpha}_{ii} &= \frac{1}{8}\left(\frac{V(\widehat{X}^{n,N}_{ii}, r, 0)^m_1}{\sqrt{\frac{1}{\pi}}\Gamma\left(\frac{r+1}{2}\right)\frac{1}{m}\sum_{p=1}^m \left(\widehat{X}^{n,N}_{t_p^m,ii}\right)^{\frac{r}{2}}} ds\right)^{\frac{2}{r}} \\
\widehat{\alpha}_{ij} &= \underset{\alpha_{12}\in[-\sqrt{\widehat{\alpha}_{11}\widehat{\alpha}_{22}},\sqrt{\widehat{\alpha}_{11}\widehat{\alpha}_{22}}]}{\operatorname{argmin}} \Big(V(\widehat{X}^{n,N}_{12}, f_r, 0)^m_1 - PV_{12}(\widehat{X}^{n,N}, \widehat{\alpha}_{11}, \alpha_{12}, \widehat{\alpha}_{22})\Big)^2 \\
\widehat{\rho} &= \underset{\rho\in[-1,1]^2,\, \rho^\top\rho\leq 1}{\operatorname{argmin}} \sum_{i=1}^{2}\Big(V(\widehat{X}^{n,N}_{ii}, Y_i, f_{r,s}, 0)^m_1 - PC_{ii}(\widehat{X}^{n,N}, Y, \widehat{\alpha}, \rho)\Big)^2.
\end{aligned}
$$

In our simulation study, we choose $r = \frac{1}{4}$ in $V(\widehat{X}^{n,N}_{11}, f_r, 0)^m_1$ and $r = 1$ in $V(\widehat{X}^{n,N}_{ij}, f_r, 0)^m_1$ for $(ij) = (12)$ and $(ij) = (22)$ respectively. This is due to the fact that X_{11} exhibits jumps and taking a lower power reduces the contribution of jumps in the power variation. In $V(\widehat{X}^{n,N}_{ii}, Y_i, f_{r,s}, 0)^m_1$, r and s are chosen to be $\frac{1}{2}$. Figure 5 and 6 show the estimated values for α and ρ as a function of the grid points m. As a consequence of Theorem 7.1, the grid corresponding to m has to be coarsened considerably with respect to the initial gridding with n points (of order $n^{\frac{1}{4}}$ or even more depending on the power used). For this reason the number of grid points shown in the graphs is rather small. Nevertheless the estimation results are good approximations of the true parameter values and can further be improved by increasing n and thus in turn also m.

References

[1] O. E. Barndorff-Nielsen, S. E. Graversen, J. Jacod, M. Podolskij, and N. Shephard. A central limit theorem for realised power and bipower variations of continuous semimartingales. In *From stochastic calculus to mathematical finance*, pages 33–68. Springer, Berlin, 2006.

[2] O. E. Barndorff-Nielsen, N. Shephard, and M. Winkel. Limit theorems for multipower variation in the presence of jumps. 2005.

[3] E. Eberlein and D. B. Madan. Sato processes and the valuation of structured products. 2011.

[4] J. Jacod. Asymptotic properties of realized power variations and related functionals of semimartingales. *Stochastic Processes and their Applications*, 118(4):517–559, 2008.

[5] J. Jacod and P. Protter. *Discretization of processes*, volume 67 of *Stochastic Modelling and Applied Probability*. Springer, Heidelberg, 2012.

[6] J. Jacod and A. N. Shiryaev. *Limit theorems for stochastic processes*, volume 288 of *Fundamental Principles of Mathematical Sciences*. Springer-Verlag, Berlin, second edition, 2003.

[7] G. G. Lorentz. *Approximation of functions*. Chelsea Publishing Co., New York, second edition, 1986.

[8] P. Malliavin and M. E. Mancino. Fourier series method for measurement of multivariate volatilities. *Finance Stoch.*, 6(1):49–61, 2002.

[9] P. Malliavin and M. E. Mancino. A Fourier transform method for nonparametric estimation of multivariate volatility. *Ann. Statist.*, 37(4):1983–2010, 2009.

[10] S. Nabeya. Absolute moments in 2-dimensional normal distribution. *Ann. Inst. Statist. Math., Tokyo*, 3:2–6, 1951.

[11] M. Podolskij. New theory on estimation of integrated volatility with applications. PhD Thesis, Ruhr-Universitaet Bochum.

[12] V. Todorov and G. Tauchen. The realized Laplace transforms of volatility. Preprint, 2011.

[13] J. H. C. Woerner. Power and multipower variation: inference for high frequency data. In A. N. Shiryaev, M. R. Grossinho, P. E. Oliveira, and M. L. Esquivel, editors, *Stochastic Finance*, pages 343–364. Springer US, 2006.



Figure 5. Estimation of α as a function of the grid points

University of Vienna, Faculty of Mathematics, Nordbergstrasse 15, A-1090 Wien, Austria,
ETH Zürich, D-Math, Rämistrasse 101, CH-8092 Zürich, Switzerland
E-mail address: `christa.cuchiero@univie.ac.at, jteichma@math.ethz.ch`



FIGURE 6. Estimation of ρ as a function of the grid points